

				<p>factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical</p>	<p>described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms</p>
--	--	--	--	---	---

				<p>approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p> <p>and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an</p>
--	--	--	--	---

228	HHEPD24	1176	Production of IL-6	<p>IL-6 FMAT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and</p>	<p>infectious disease as described below under "Infectious Disease").</p> <p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as</p>
-----	---------	------	--------------------	--	--

				<p>differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using</p>	<p>described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and</p>
--	--	--	--	--	---

				<p>techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
228	HHEPD24	1176	Production of MCP-1	<p>MCP-1 FMAT. Assays for immunomodulatory proteins that are produced by a large variety of cells and act to induce chemotaxis and activation of monocytes and T</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) MCP-1 production. An alternative highly preferred</p>

			<p>cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, induce chemotaxis, and modulate immune cell activation. Exemplary assays that test for immunomodulatory proteins evaluate the production of cell surface markers, such as monocyte chemoattractant protein (MCP), and the activation of monocytes and T cells. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Satthaporn and</p>	<p>embodiment of the invention includes a method for inhibiting (e.g., reducing) MCP-1 production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL),</p>
--	--	--	---	--

				<p>Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety.</p> <p>Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art.</p> <p>Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis (bacterial and viral), Lyme Disease, asthma, and allergy Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia,</p>
--	--	--	--	--	--

228	HHEPD24	1176	Production of MIP1alpha	<p>MIP-1alpha FMAT. Assays for immunomodulatory proteins produced by activated dendritic cells that upregulate monocyte/macrophage and T cell chemotaxis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate chemotaxis, and modulate T cell differentiation. Exemplary assays that test for immunomodulatory proteins evaluate the production of chemokines, such as macrophage inflammatory protein 1 alpha (MIP-1a), and the activation of monocytes/macrophages and T cells. Such assays that may be used or routinely modified to test immunomodulatory and chemotaxis activity of polypeptides of the invention (including antibodies and</p>	<p>metaplasia, and/or dysplasia.</p> <p>A highly preferred embodiment of the invention includes a method for stimulating MIP1a production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MIP1a production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders.</p>
-----	---------	------	-------------------------	--	--

				<p>agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); Drakes et al., Transp Immunol 8(1):17-29 (2000); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma, and allergy. Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include</p>
--	--	--	--	--	---

					benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
228	HHEPD24	1176	Production of TNF alpha by dendritic cells	<p>TNFα F/MAT. Assays for immunomodulatory proteins produced by activated macrophages, T cells, fibroblasts, smooth muscle, and other cell types that exert a wide variety of inflammatory and cytotoxic effects on a variety of cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate inflammation and cytotoxicity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines such as tumor necrosis factor alpha (TNFα), and the induction or inhibition of an inflammatory or cytotoxic response. Such</p>	<p>A highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and</p>

				<p>assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Verhasselt et al., Eur J Immunol 28(11):3886-3890 (1198); Dahlen et al., J Immunol 160(7):3585-3593 (1998); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which,</p>	<p>suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include</p>
--	--	--	--	--	--

					when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.	anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
229	HHEPM33	1177	SEAP in 293/ISRE			
229	HHEPM33	1177	Activation of transcription through cAMP response element (CRE) in pre-adipocytes.	Assays for the activation of transcription through the cAMP response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention	A highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. An additional highly preferred	

				<p>(including antibodies and agonists or antagonists of the invention) to increase cAMP, regulate CREB transcription factors, and modulate expression of genes involved in a wide variety of cell functions. For example, a 3T3-L1/CRE reporter assay may be used to identify factors that activate the cAMP signaling pathway. CREB plays a major role in adipogenesis, and is involved in differentiation into adipocytes. CRE contains the binding sequence for the transcription factor CREB (CRE binding protein). Exemplary assays for transcription through the cAMP response element that may be used or routinely modified to test cAMP-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and</p>	<p>indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hypermolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as</p>
--	--	--	--	--	--

				<p>Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Reusch et al., Mol Cell Biol 20(3):1008-1020 (2000); and Klemm et al., J Biol Chem 273:917-923 (1998), the contents of each of which are herein incorporated by reference in its entirety. Pre-adipocytes that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). Additional highly preferred indications are complications associated with insulin resistance.</p>
229	HHEPM33	1177	Activation of transcription	<p>This reporter assay measures activation of the GATA-3</p>	<p>Highly preferred indications include allergy, asthma, and</p>

			through GATA-3 response element in immune cells (such as mast cells).	<p>signaling pathway in HMC-1 human mast cell line. Activation of GATA-3 in mast cells has been linked to cytokine and chemokine production. Assays for the activation of transcription through the GATA3 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate GATA3 transcription factors and modulate expression of mast cell genes important for immune response development. Exemplary assays for transcription through the GATA3 response element that may be used or routinely modified to test GATA3-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and</p>	<p>rhinitis. Additional preferred indications include infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications also include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary tract cancers and/or as described below under "Hyperproliferative Disorders"). Other preferred</p>
--	--	--	---	--	--

				<p>Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Flavell et al., Cold Spring Harb Symp Quant Biol 64:563-571 (1999); Rodriguez-Palmero et al., Eur J Immunol 29(12):3914-3924 (1999); Zheng and Flavell, Cell 89(4):587-596 (1997); and Henderson et al., Mol Cell Biol 14(6):4286-4294 (1994), the contents of each of which are herein incorporated by reference in its entirety. Mast cells that may be used according to these assays are publicly available (e.g., through the ATCC).</p> <p>Exemplary human mast cells that may be used according to these assays include the HMC-1 cell line, which is an immature human mast cell line established from the peripheral blood of a patient with mast cell leukemia, and exhibits many characteristics of immature mast cells.</p>	<p>indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.</p>
229	HHEPM33	1177	Activation of transcription	<p>This reporter assay measures activation of the NFAT</p>	<p>Highly preferred indications include allergy, asthma, and</p>

			<p>through NFAT response element in immune cells (such as mast cells).</p>	<p>signaling pathway in HMC-1 human mast cell line. Activation of NFAT in mast cells has been linked to cytokine and chemokine production. Assays for the activation of transcription through the Nuclear Factor of Activated T cells (NFAT) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFAT transcription factors and modulate expression of genes involved in immunomodulatory functions. Exemplary assays for transcription through the NFAT response element that may be used or routinely modified to test NFAT-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays</p>	<p>rhinitis. Additional preferred indications include infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications also include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary tract cancers and/or as described below under "Hyperproliferative Disorders"). Other preferred</p>
--	--	--	--	---	--

				<p>disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); De Boer et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Ali et al., J Immunol 165(12):7215-7223 (2000); Hutchinson and McCloskey, J Biol Chem 270(27):16333-16338 (1995), and Turner et al., J Exp Med 188:527-537 (1998), the contents of each of which are herein incorporated by reference in its entirety. Mast cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human mast cells that may be used according to these assays include the HMC-1 cell line, which is an immature human mast cell line established from the peripheral blood of a patient with mast cell leukemia, and exhibits many characteristics of immature mast cells.</p>	<p>indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.</p>
--	--	--	--	--	--

229	HHEPM33	1177	<p>Activation of transcription through NFAT response element in immune cells (such as natural killer cells).</p>	<p>Assays for the activation of transcription through the Nuclear Factor of Activated T cells (NFAT) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFAT transcription factors and modulate expression of genes involved in immunomodulatory functions. Exemplary assays for transcription through the NFAT response element that may be used or routinely modified to test NFAT-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988);</p>	<p>Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative</p>
-----	---------	------	--	---	--

				<p>Aramburu et al., J Exp Med 182(3):801-810 (1995); De Boer et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Fraser et al., Eur J Immunol 29(3):838-844 (1999); and Yeseen et al., J Biol Chem 268(19):14285-14293 (1993), the contents of each of which are herein incorporated by reference in its entirety. NK cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human NK cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and cytotoxic activity.</p>	<p>Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
--	--	--	--	--	---

229	HHPEM33	1177	<p>Activation of transcription through serum response element in immune cells (such as natural killer cells).</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-related genes in many cell types. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and</p>
-----	---------	------	---	---	--

				<p>3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and cytotoxic activity.</p>	<p>treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL),</p>
--	--	--	--	---	--

					<p>plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
229	HHEPM33	1177	SEAP in NK16/STAT6		
229	HHEPM33	1177	Hexosaminidase in RBL-2H3		
229	HHEPM33	1177	Activation of transcription through GAS response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies	Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma

			<p>and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Hentinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary human T cells, such as the SUPT cell line, that may be used according to these</p>	<p>(e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under</p>
--	--	--	--	---

					<p>assays are publicly available (e.g., through the ATCC).</p>	<p>“Immune Activity”, “Blood-Related Disorders”, and/or “Cardiovascular Disorders”), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under “Infectious Disease”). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
--	--	--	--	--	--	--

230	HHPT60	1178	<p>Activation of transcription through NFKB response element in immune cells (such as natural killer cells).</p>	<p>Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Valle Blazquez et al, Immunology 90(3):455-460 (1997);</p>	<p>Highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), and immunodeficiencies (e.g., as described below). An additional highly preferred indication is infection (e.g., AIDS, and/or an infectious disease as described below under "Infectious Disease"). Highly preferred indications include neoplastic diseases (e.g., melanoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for</p>
-----	--------	------	--	--	--

				<p>Aramburau et al., J Exp Med 82(3):801-810 (1995); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. NK cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human NK cells that may be used according to these assays include the NKL cell line, which is a human natural killer cell line established from the peripheral blood of a patient with large granular lymphocytic leukemia. This IL-2 dependent suspension culture cell line has a morphology resembling that of activated NK cells.</p>	<p>example, melanoma, renal cell carcinoma, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, suppression of immune reactions to transplanted organs, asthma and allergy.</p>
231	HHEPU04	1179	Production of IL-6	IL-6 FMAT. IL-6 is produced by T cells and has strong	A highly preferred embodiment of the invention

				<p>effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of</p>	<p>includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a</p>
--	--	--	--	--	--

			<p>cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhaselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which,</p>	<p>B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia,</p>
--	--	--	---	--

				when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.	leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
231	HHEPU04	1179	Production of TNF alpha by dendritic cells	TNFα FMAT. Assays for immunomodulatory proteins produced by activated macrophages, T cells, fibroblasts, smooth muscle, and other cell types that exert a wide variety of inflammatory and cytotoxic effects on a variety of cells are well known in the art and may be used or routinely modified to assess	A highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Highly preferred indications

				<p>the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate inflammation and cytotoxicity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines such as tumor necrosis factor alpha (TNFa), and the induction or inhibition of an inflammatory or cytotoxic response. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Verhasselt et al., Eur J Immunol 28(11):3886-3890 (1198); Dahlen et al., J Immunol 160(7):3585-3593</p>	<p>include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications</p>
--	--	--	--	---	---

				<p>(1998); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease,</p>
--	--	--	--	--	---

					cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
232	HHFBY53	1180	Activation of Endothelial Cell p38 or JNK Signaling Pathway.	<p>Kinase assay. JNK and p38 kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for</p>

			<p>Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are endothelial cells which line venous blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) endothelial cell activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) the activation of and/or inactivating endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as</p>
--	--	--	--	--

					described below under “Hyperproliferative Disorders”), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under “Cardiovascular Disorders”). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly
--	--	--	--	--	--

					<p>preferred are indications that inhibit angiogenesis and/or cardiovascularization.</p> <p>Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders.</p> <p>Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
--	--	--	--	--	--

					<p>Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud"s disease and Reynaud"s phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke,</p>
--	--	--	--	--	--

					<p>graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment /prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease.</p> <p>Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders").</p> <p>Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such</p>
--	--	--	--	--	--

					as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.
233	HHFEC49	1181	Production of IFNgamma using a T cells	<p>IFNgamma FMAT. IFNγ plays a central role in the immune system and is considered to be a proinflammatory cytokine. IFNγ promotes TH1 and inhibits TH2 differentiation; promotes IgG2a and inhibits IgE secretion; induces macrophage activation; and increases MHC expression. Assays for immunomodulatory proteins produced by T cells and NK cells that regulate a variety of inflammatory activities and inhibit TH2 helper cell functions are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating the production of IFNγ. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of IFNγ. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic</p>

				<p>function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon gamma (IFNg), and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated by</p>	<p>lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiency (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic</p>
--	--	--	--	---	--

				reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.	conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.
234	HHFFJ48	1182	Activation of Adipocyte PI3 Kinase Signalling Pathway	Kinase assay. Kinase assays, for example an GSK-3 assays, for PI3 kinase signal transduction that regulate glucose metabolism and cell survival are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including	<p>A highly preferred embodiment of the invention includes a method for increasing adipocyte survival</p> <p>An alternative highly preferred embodiment of the invention includes a method for decreasing adipocyte survival.</p> <p>A preferred embodiment of the invention includes a method</p>

			<p>antibodies and agonists or antagonists of the invention) to promote or inhibit glucose metabolism and cell survival. Exemplary assays for PI3 kinase activity that may be used or routinely modified to test PI3 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Nikoulina et al., Diabetes 49(2):263-271 (2000); and Schreyer et al., Diabetes 48(8):1662-1666 (1999), the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a</p>	<p>for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"). Preferred indications include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"), blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-</p>
--	--	--	---	---

				<p>continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>Related Disorders”), immune disorders (e.g., as described below under “Immune Activity”), neural disorders (e.g., as described below under “Neural Activity and Neurological Diseases”), and infection (e.g., as described below under “Infectious Disease”). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the “Renal Disorders” section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness,</p>
--	--	--	--	--	--

					<p>nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional</p>
--	--	--	--	--	--

					<p>highly preferred indications are complications associated with insulin resistance.</p> <p>Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein.</p> <p>Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Highly preferred indications include neoplasms and cancer, such as, lipoma, liposarcoma, lymphoma, leukemia and breast, colon, and kidney cancer. Additional highly preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such</p>
--	--	--	--	--	---

					as, for example, hyperplasia, metaplasia, and/or dysplasia.
235	HHFGR93	1183	SEAP in Alk Phos C2C12		
235	HHFGR93	1183	Production of MCP-1	MCP-1 FMAT. Assays for immunomodulatory proteins that are produced by a large variety of cells and act to induce chemotaxis and activation of monocytes and T cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, induce chemotaxis, and modulate immune cell activation. Exemplary assays that test for immunomodulatory proteins evaluate the production of cell surface markers, such as monocyte chemoattractant protein (MCP), and the activation of monocytes and T cells. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of	A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) MCP-1 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MCP-1 production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis,

			<p>polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis (bacterial and viral), Lyme Disease, asthma, and allergy Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms</p>
--	--	--	---	---

					and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
235	HHFGR93	1183	Activation of transcription through NFKB response element in epithelial cells (such as HELA cells).	Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of epithelial genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides of the invention (including antibodies	Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Cancer, Wound Healing, and Inflammation. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and

				and agonists or antagonists of the invention) include assays disclosed in: Kaltschmidt B, et al., <i>Oncogene</i> , 18(21):3213-3225 (1999); Beetz A, et al., <i>Int J Radiat Biol</i> , 76(11):1443-1453 (2000); Berger et al., <i>Gene</i> 66:1-10 (1998); Cullen and Malm, <i>Methods in Enzymol</i> 216:362-368 (1992); Henthorn et al., <i>Proc Natl Acad Sci USA</i> 85:6342-6346 (1988); Valle Blazquez et al., <i>Immunology</i> 90(3):455-460 (1997); Aramburau et al., <i>J Exp Med</i> 82(3):801-810 (1995); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Epithelial cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary epithelial cells that may be used according to these assays include the HELA cell line.	urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include inflammation and inflammatory disorders.
	HHFGR93	1183	IFNg in Human T-cell 2B9		

235	HHFGR93	1183	Production of RANTES in bronchial epithelium cells	<p>RANTES FMAT. Assays for immunomodulatory proteins that induce chemotaxis of T cells, monocytes, and eosinophils are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, induce chemotaxis, and/or mediate humoral or cell-mediated immunity.</p> <p>Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as RANTES, and the induction of chemotactic responses in immune cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-</p>	
-----	---------	------	--	--	--

				204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Cocchi et al., Science 270(5243):1811-1815 (1995); and Robinson et al., Clin Exp Immunol 101(3):398-407 (1995), the contents of each of which are herein incorporated by reference in its entirety. Epithelial cells were isolated from bronchia/trachea immediately postmortem from humans who were free of known respiratory diseases. See Wu et al., Am Rev Respir Dis. 132(2):311-20 (1985), the contents of which are herein incorporated by reference in its entirety.	
235	HHFGR93	1183	Calcium flux in immune cells (such as monocytes)	Assays for measuring calcium flux are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mobilize calcium. Cells normally have very low concentrations of cytosolic calcium compared to	Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Infection, Inflammation, Atherosclerosis, Hypersensitivity, and Leukemias

				<p>much higher extracellular calcium. Extracellular factors can cause an influx of calcium, leading to activation of calcium responsive signaling pathways and alterations in cell functions. Exemplary assays that may be used or routinely modified to measure calcium flux in immune cells (such as monocytes) include assays disclosed in: Chan, CC, et al., J Pharmacol Exp Ther, 269(3):891-896 (1994); Andersson, K, et al., Cytokine, 12(12):1784-1787 (2000); Scully, SP, et al., J Clin Invest, 74(2) 589-599 (1984); and, Sullivan, E, et al., Methods Mol Biol, 114:125-133 (1999), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include the THP-1 monocyte</p>	
--	--	--	--	--	--

236	HHFHJ59	1184	Production of IL-10 and activation of T-cells.	cell line. Assays for production of IL-10 and activation of T-cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate or inhibit production of IL-10 and/or activation of T-cells. Exemplary assays that may be used or routinely modified to assess the ability of polypeptides and antibodies of the invention (including agonists or antagonists of the invention) to modulate IL-10 production and/or T-cell proliferation include, for example, assays such as disclosed and/or cited in: Robinson, DS, et al., "Th-2 cytokines in allergic disease" Br Med Bull; 56 (4): 956-968 (2000), and Cohn, et al., "T-helper type 2 cell-directed therapy for asthma" Pharmacology & Therapeutics; 88: 187-196 (2000); the	Highly preferred indications include allergy and asthma. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response.
-----	---------	------	--	---	---

				contents of each of which are herein incorporated by reference in their entirety. Exemplary cells that may be used according to these assays include Th2 cells. IL10 secreted from Th2 cells may be measured as a marker of Th2 cell activation. Th2 cells are a class of T cells that secrete IL4, IL10, IL13, IL5 and IL6. Factors that induce differentiation and activation of Th2 cells play a major role in the initiation and pathogenesis of allergy and asthma. Primary T helper 2 cells are generated via in vitro culture under Th2 polarizing conditions using peripheral blood lymphocytes isolated from cord blood.	
237	HHFHR32	1185	Production of IL-6	IL-6 FMAT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6	A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A

			<p>has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test</p>	<p>highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include</p>
--	--	--	---	--

				<p>immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease,</p>
--	--	--	--	--	--

					sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
238	HHFOJ29	1186	SEAP in Jurkat/IL4 promoter (antiCD3 co-stim)		1
238	HHFOJ29	1186	Activation of Natural Killer Cell ERK Signaling Pathway.	Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK	A highly preferred embodiment of the invention includes a method for stimulating natural killer cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting natural killer cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating natural killer cell differentiation. An alternative highly preferred embodiment of the invention

				<p>kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Natural killer cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary natural killer cells that may be used according to these assays include the human natural killer cell lines (for example, NK-YT cells which have cytolytic and cytotoxic activity) or primary NK cells.</p>	<p>includes a method for inhibiting natural killer cell differentiation. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity") and infections (e.g., as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as</p>
--	--	--	--	---	---

					described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include cancers such as, kidney, melanoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, urinary cancer, lymphoma and leukemias. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Other highly preferred indications include, pancytopenia, leukopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), arthritis, asthma, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, immune reactions to transplanted organs and tissues, endocarditis, meningitis, Lyme Disease, and allergies.
	HHGB091	1187	Activation of	Kinase assay. Kinase assays,	A highly preferred

239	Adipocyte ERK Signaling Pathway	<p>for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and</p>	<p>embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A highly preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) adipocyte activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of (e.g., decreasing) and/or inactivating adipocytes. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders").</p>
-----	---------------------------------	--	--

				<p>Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety.</p> <p>Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC).</p> <p>Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>Highly preferred indications also include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease").</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic</p>
--	--	--	--	---	---

					<p>nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hypermolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired</p>
--	--	--	--	--	---

					<p>wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below (particularly of the urinary tract and skin). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p> <p>Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein.</p> <p>Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Preferred</p>
--	--	--	--	--	---

					indications include neoplasms and cancer, such as, lymphoma, leukemia and breast, colon, and kidney cancer. Additional preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Highly preferred indications include lipomas and liposarcomas. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
240	HHGCM76	1188	IL-10 in Human T-cell 293T		
240	HHGCM76	1188	Stimulation of insulin secretion from pancreatic beta cells.	Assays for measuring secretion of insulin are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin secretion. For example, insulin secretion is measured by FMAT using anti-rat insulin antibodies.	A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below),

			<p>Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain proteins/peptides, and dysregulation is a key component in diabetes. Exemplary assays that may be used or routinely modified to test for stimulation of insulin secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Ahren, B., et al., Am J Physiol, 277(4 Pt 2):R959-66 (1999); Li, M., et al., Endocrinology, 138(9):3735-40 (1997); Kim, K.H., et al., FEBS Lett, 377(2):237-9 (1995); and, Miraglia S et. al., Journal of Biomolecular Screening, 4:193-204 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC)</p>	<p>diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal</p>
--	--	--	--	--

			<p>and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an X-ray induced rat transplantable insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Asfari et al. Endocrinology 1992 130:167.</p>	<p>tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
240	HHGCM76	1188	<p>Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Takacs P, et al, FASEB J, 15(2):279-281</p>	<p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Inflammation, Vascular Disease, Atherosclerosis, Restenosis, and Stroke</p>

241	HHGCQ54	1189	Activation of Adipocyte ERK Signaling Pathway	<p>(2001); and, Miyamoto K, et al., Am J Pathol, 156(5):1733-1739 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include microvascular endothelial cells (MVEC).</p> <p>Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A highly preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte</p>
-----	---------	------	---	---	---

				<p>test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3</p>	<p>differentiation. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) adipocyte activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of (e.g., decreasing) and/or inactivating adipocytes. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"). Highly preferred indications also include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related</p>
--	--	--	--	---	---

				<p>fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease").</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-</p>
--	--	--	--	--	---

					<p>hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below (particularly of the urinary tract and skin). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
--	--	--	--	--	---

					<p>Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein.</p> <p>Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Preferred indications include neoplasms and cancer, such as, lymphoma, leukemia and breast, colon, and kidney cancer. Additional preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer.</p> <p>Highly preferred indications include lipomas and liposarcomas. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia,</p>
--	--	--	--	--	--

242	HHGDF16	1190	Activation of transcription through serum response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the	metaplasia, and/or dysplasia. A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and
-----	---------	------	---	---	---

				<p>content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute</p>
--	--	--	--	--	--

					lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
242	HHGDF16	1190	Activation of transcription through AP1 response element in immune cells (such as T-cells).	Assays for the activation of transcription through the AP1 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for	Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under

				<p>transcription through the AP1 response element that may be used or routinely modified to test AP1-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., <i>Gene</i> 66:1-10 (1988); Cullen and Malm, <i>Methods in Enzymol</i> 216:362-368 (1992); Henthorn et al., <i>Proc Natl Acad Sci USA</i> 85:6342-6346 (1988); Rellahan et al., <i>J Biol Chem</i> 272(49):30806-30811 (1997); Chang et al., <i>Mol Cell Biol</i> 18(9):4986-4993 (1998); and Fraser et al., <i>Eur J Immunol</i> 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is an IL-2 and IL-4</p>	<p>“Infectious Disease”). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under “Hyperproliferative Disorders”). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
--	--	--	--	---	--

				responsive suspension-culture cell line.	Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.
243	HHGDW43	1191	Activation of transcription through API response element in immune cells (such as T-cells).	Assays for the activation of transcription through the API response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for transcription through the API response element that may be used or routinely modified to test API-response element	Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic

				<p>activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is an IL-2 and IL-4 responsive suspension-culture cell line.</p>	<p>lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia,</p>
--	--	--	--	---	--

243	HHGDW43	1191	Production of IL-10 and activation of T-cells.	Assays for production of IL-10 and activation of T-cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate or inhibit production of IL-10 and/or activation of T-cells. Exemplary assays that may be used or routinely modified to assess the ability of polypeptides and antibodies of the invention (including agonists or antagonists of the invention) to modulate IL-10 production and/or T-cell proliferation include, for	Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.
				Assays for production of IL-10 and activation of T-cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate or inhibit production of IL-10 and/or activation of T-cells. Exemplary assays that may be used or routinely modified to assess the ability of polypeptides and antibodies of the invention (including agonists or antagonists of the invention) to modulate IL-10 production and/or T-cell proliferation include, for	Highly preferred indications include allergy and asthma. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response.

				<p>example, assays such as disclosed and/or cited in: Robinson, DS, et al., "Th-2 cytokines in allergic disease" Br Med Bull; 56 (4): 956-968 (2000), and Cohn, et al., "T-helper type 2 cell-directed therapy for asthma" Pharmacology & Therapeutics; 88: 187-196 (2000); the contents of each of which are herein incorporated by reference in their entirety. Exemplary cells that may be used according to these assays include Th2 cells. IL10 secreted from Th2 cells may be measured as a marker of Th2 cell activation. Th2 cells are a class of T cells that secrete IL4, IL10, IL13, IL5 and IL6. Factors that induce differentiation and activation of Th2 cells play a major role in the initiation and pathogenesis of allergy and asthma. Primary T helper 2 cells are generated via in vitro culture under Th2 polarizing conditions using peripheral blood lymphocytes isolated</p>
--	--	--	--	--

244	HHPDX20	1192	<p>Activation of transcription through cAMP response element (CRE) in pre-adipocytes.</p>	<p>from cord blood.</p> <p>Assays for the activation of transcription through the cAMP response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to increase cAMP, regulate CREB transcription factors, and modulate expression of genes involved in a wide variety of cell functions. For example, a 3T3-L1/CRE reporter assay may be used to identify factors that activate the cAMP signaling pathway. CREB plays a major role in adipogenesis, and is involved in differentiation into adipocytes. CRE contains the binding sequence for the transcription factor CREB (CRE binding protein). Exemplary assays for transcription through the cAMP response element that may be used or routinely</p>	<p>A highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. An additional highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g.,</p>
-----	---------	------	---	--	--

				<p>modified to test cAMP-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Reusch et al., Mol Cell Biol 20(3):1008-1020 (2000); and Klemm et al., J Biol Chem 273:917-923 (1998), the contents of each of which are herein incorporated by reference in its entirety. Preadipocytes that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3</p>	<p>heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). Additional highly preferred indications are complications associated with insulin resistance.</p>
--	--	--	--	---	---

244	HHPDX20	1192	Endothelial Cell Apoptosis	<p>fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p> <p>Caspase Apoptosis. Assays for caspase apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote caspase protease-mediated apoptosis. Induction of apoptosis in endothelial cells supporting the vasculature of tumors is associated with tumor regression due to loss of tumor blood supply. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Lee et al., FEBS</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for</p>
-----	---------	------	----------------------------	---	---

				<p>inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular</p>
			<p>Lett 485(2-3): 122-126 (2000); Nor et al., J Vasc Res 37(3): 209-218 (2000); and Karsan and Harlan, J Atheroscler Thromb 3(2): 75-80 (1996); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through commercial sources). Exemplary endothelial cells that may be used according to these assays include bovine aortic endothelial cells (bAEC), which are an example of endothelial cells which line blood vessels and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	

					<p>disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders").</p> <p>Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization.</p> <p>Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma,</p>
--	--	--	--	--	---

hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis,					
---	--	--	--	--	--

					<p>lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment /prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas,</p>
--	--	--	--	--	---

					heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.
244	HHPDX20	1192	IL-10 in Human T-cell 293T		
245	HHPGO40	1193	Proliferation of immune cells (such as the HMC-1 human mast cell	Assays for the regulation (i.e. increases or decreases) of viability and proliferation of cells in vitro are well-known in	Highly preferred indications include asthma, allergy, mastocytosis (a rare, heterogeneous disorder

			line)	<p>the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate viability and proliferation of eosinophil cells and cell lines. For example, the CellTiter-Glo[®] Luminescent Cell Viability Assay (Promega Corp., Madison, WI, USA) can be used to measure the number of viable cells in culture based on quantitation of the ATP present which signals the presence of metabolically active cells. Mast cells are found in connective and mucosal tissues throughout the body. Mast cell activation (via immunoglobulin E -antigen, promoted by T helper cell type 2 cytokines) is an important component of allergic disease. Dysregulation of mast cell apoptosis may play a role in allergic disease and mast cell tumor survival. Mast cell lines that may be used according to</p>	<p>characterized by excessive accumulation of mast cells, and their proliferation and action in the skin, central nervous system, and other organs). Preferred indications also include hematopoietic and immunological disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), infection (e.g., as described below under "Infectious Disease"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), and immunodeficiencies (e.g., as described below).</p>
--	--	--	-------	---	--

				these assays are publicly available and/or may be routinely generated. Exemplary mast cells that may be used according to these assays include HMC-1, which is an immature human mast cell line established from the peripheral blood of a patient with mast cell leukemia, and exhibits many characteristics of immature mast cells.	
245	HHPGO40	1193	Glucose Production in H4IIE		
245	HHPGO40	1193	Proliferation of pre-adipose cells (such as 3T3-L1 cells)	Assays for the regulation (i.e. increases or decreases) of viability and proliferation of cells in vitro are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate viability and proliferation of pre-adipose cells and cell lines. For example, the CellTiter-Glo [®] Luminescent Cell Viability Assay (Promega Corp., Madison, WI, USA) can be	

				used to measure the number of viable cells in culture based on quantitation of the ATP present which signals the presence of metabolically active cells. 3T3-L1 is a mouse preadipocyte cell line. It is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation. Cells were differentiated to an adipose-like state before being used in the screen. See Green H and Meuth M., Cell 3: 127-133 (1974), which is herein incorporated by reference in its entirety.	
245	HHPGO40	1193	Caspase (+pachitaxel) in SW480		
246	HHPTJ65	1194	Regulation of apoptosis of immune cells (such as mast cells).	Caspase Apoptosis. Assays for caspase apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate caspase protease-mediated apoptosis in immune cells (such as, for	Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of asthma, allergy, hypersensitivity and inflammation.

				<p>example, in mast cells). Mast cells are found in connective and mucosal tissues throughout the body, and their activation via immunoglobulin E - antigen, promoted by T helper cell type 2 cytokines, is an important component of allergic disease. Dysregulation of mast cell apoptosis may play a role in allergic disease and mast cell tumor survival. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis activity induced by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in: Masuda A, et al., <i>J Biol Chem</i>, 276(28):26107-26113 (2001); Yeatman CF 2nd, et al., <i>J Exp Med</i>, 192(8):1093-1103 (2000); Lee et al., <i>FEBS Lett</i> 485(2-3): 122-126 (2000); Nor et al., <i>J Vasc Res</i> 37(3): 209-218 (2000); and Karsan and Harlan, <i>J Atheroscler Thromb</i> 3(2): 75-80 (1996); the</p>
--	--	--	--	---

				<p>contents of each of which are herein incorporated by reference in its entirety. Immune cells that may be used according to these assays are publicly available (e.g., through commercial sources). Exemplary immune cells that may be used according to these assays include mast cells such as the HMC human mast cell line.</p>	
246	HHPTJ65	1194	IL-2 in Human T-cell 2B9		
247	HHSDX28	1195	Activation of transcription through serum response element in immune cells (such as T-cells).	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications</p>

				<p>activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast,</p>
--	--	--	--	---	--

lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below					
---	--	--	--	--	--

247	HHSDX28	1195	Production of TNF alpha by dendritic cells	<p>TNFα FMT. Assays for immunomodulatory proteins produced by activated macrophages, T cells, fibroblasts, smooth muscle, and other cell types that exert a wide variety of inflammatory and cytotoxic effects on a variety of cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate inflammation and cytotoxicity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines such as tumor necrosis factor alpha (TNFα), and the induction or inhibition of an inflammatory or cytotoxic response. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention</p>	<p>under "Infectious Disease").</p> <p>A highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and</p>
-----	---------	------	--	--	--

			<p>(including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Verhasselt et al., Eur J Immunol 28(11):3886-3890 (1998); Dahlen et al., J Immunol 160(7):3585-3593 (1998); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL),</p>
--	--	--	---	---

					<p>plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
247	HHSDX28	1195	<p>Activation of transcription through GATA-3 response element in immune cells (such as mast cells).</p>	<p>This reporter assay measures activation of the GATA-3 signaling pathway in HMC-1 human mast cell line. Activation of GATA-3 in mast cells has been linked to cytokine and chemokine production. Assays for the activation of transcription through the GATA3 response element are well-known in the art and may be used or routinely modified to assess</p>	<p>Highly preferred indications include allergy, asthma, and rhinitis. Additional preferred indications include infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications also include blood disorders (e.g., as described below under "Immune Activity", "Blood-</p>

			<p>the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate GATA3 transcription factors and modulate expression of mast cell genes important for immune response development. Exemplary assays for transcription through the GATA3 response element that may be used or routinely modified to test GATA3-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Flavell et al., Cold Spring Harb Symp Quant Biol 64:563-571 (1999); Rodriguez-Palmero et al., Eur J Immunol 29(12):3914-3924 (1999); Zheng and Flavell, Cell 89(4):587-596 (1997); and Henderson et al., Mol Cell Biol</p>	<p>Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary tract cancers and/or as described below under "Hyperproliferative Disorders"). Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas,</p>
--	--	--	---	---

				<p>14(6):4286-4294 (1994), the contents of each of which are herein incorporated by reference in its entirety. Mast cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human mast cells that may be used according to these assays include the HMC-1 cell line, which is an immature human mast cell line established from the peripheral blood of a patient with mast cell leukemia, and exhibits many characteristics of immature mast cells.</p>	<p>multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.</p>
247	HHSDX28	1195	IgG in Human B cells		
247	HHSDX28	1195	MCP-1 in HUVEC		
248	HILCF66	1196	TNFa in Human T-cell 293T		
248	HILCF66	1196	Activation of Endothelial Cell p38 or JNK Signaling Pathway.	<p>Kinase assay. JNK and p38 kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method</p>

			<p>the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g.,</p>	<p>for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) endothelial cell activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) the activation of and/or inactivating endothelial cells.</p>
--	--	--	---	---

				<p>through the ATCC). Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are endothelial cells which line venous blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac</p>
--	--	--	--	--	---

					<p>hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders"). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization. Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors,</p>
--	--	--	--	--	---

					<p>telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other</p>
--	--	--	--	--	---

vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment /prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and					
--	--	--	--	--	--

					<p>vascular disease.</p> <p>Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders").</p> <p>Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.</p>
249	HJACG02	1197	MIP-1a in HMC		
249	HJACG02	1197	Proliferation of pre-adipose cells (such as 3T3-L1 cells)	Assays for the regulation (i.e. increases or decreases) of viability and proliferation of cells in vitro are well-known in the art and may be used or routinely modified to assess	

					the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate viability and proliferation of pre-adipose cells and cell lines. For example, the CellTiter-Glo [®] Luminescent Cell Viability Assay (Promega Corp., Madison, WI, USA) can be used to measure the number of viable cells in culture based on quantitation of the ATP present which signals the presence of metabolically active cells. 3T3-L1 is a mouse preadipocyte cell line. It is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation. Cells were differentiated to an adipose-like state before being used in the screen. See Green H and Meuth M., Cell 3: 127-133 (1974), which is herein incorporated by reference in its entirety.	
249	HJACG02	1197	MCP-1 in HUVEC		This reporter assay measures	
	HJACG02	1197	Activation or			

249				<p>inhibition of transcription through NFkB response element in immune cells (such as basophils).</p> <p>activation or inhibition of the NFkB signaling pathway in Ku812 human basophil cell line. Assays for the activation or inhibition of transcription through the NFkB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFkB transcription factors and modulate expression of immunomodulatory genes. NFkB is important in the pathogenesis of asthma. Exemplary assays for transcription through the NFkB response element that may be used or routinely modified to test NFkB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and</p>	
-----	--	--	--	--	--

				<p>Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Marone et al, Int Arch Allergy Immunol 114(3):207-17 (1997), the contents of each of which are herein incorporated by reference in its entirety. Cells were pretreated with SID supernatants or controls for 15-18 hours, and then 10 ng/mL of TNF was added to stimulate the NFkB reporter. SEAP activity was measured after 48 hours. Basophils that may be used according to these assays are publicly available (e.g., through the ATCC).</p> <p>Exemplary human basophil cell lines that may be used according to these assays include Ku812, originally established from a patient with chronic myelogenous leukemia. It is an immature prebasophilic cell line that can be induced to differentiate into mature basophils. See, Kishi et al., Leuk Res. 9:381-390 (1985); Blom et al., Eur J</p>
--	--	--	--	--

250	HJACG30	1198	Activation of transcription through serum response element in immune cells (such as T-cells).	Immunol. 22:2025-32 (1992), where the contents of each are herein incorporated by reference in its entirety. Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated
-----	---------	------	---	--	--

				<p>85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include</p>
--	--	--	--	---	---

					<p>anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
250	HJACG30	1198	Inhibition of squalene synthetase gene transcription.	<p>Reporter Assay: construct contains regulatory and coding sequence of squalene synthetase, the first specific enzyme in the cholesterol biosynthetic pathway. See Jiang, et al., J. Biol. Chem. 268:12818-12824(1993), the contents of which are herein</p>	

250	HJACG30	1198	Stimulation of insulin secretion from pancreatic beta cells.	<p>incorporated by reference in its entirety. Cells were treated with SID supernatants, and SEAP activity was measured after 72 hours. HepG2 is a human hepatocellular carcinoma cell line (ATCC HB-8065). See Knowles et al., Science. 209:497-9 (1980), the contents of which are herein incorporated by reference in its entirety.</p> <p>Assays for measuring secretion of insulin are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin secretion. For example, insulin secretion is measured by FMAT using anti-rat insulin antibodies. Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain proteins/peptides, and dysregulation is a key component in diabetes.</p>	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic neuropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel</p>
-----	---------	------	--	---	--

				<p>Exemplary assays that may be used or routinely modified to test for stimulation of insulin secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Ahren, B., et al., Am J Physiol, 277(4 Pt 2):R959-66 (1999); Li, M., et al., Endocrinology, 138(9):3735-40 (1997); Kim, K.H., et al., FEBS Lett, 377(2):237-9 (1995); and, Miraglia S et. al., Journal of Biomolecular Screening, 4:193-204 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established from cells</p>	<p>blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyposmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include</p>
--	--	--	--	---	---

				isolated from an X-ray induced rat transplantable insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Asfari et al. Endocrinology 1992 130:167.	weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.
251	HJBCU04	1199	Activation of transcription through GAS response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies	Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia,

				<p>and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety.</p> <p>Exemplary mouse T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the CTLL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.</p>	<p>metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders.</p> <p>Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is</p>
--	--	--	--	---	---

251	HJBCU04	1199	Production of IL-4	IL-4 FMAT. Assays for immunomodulatory proteins secreted by TH2 cells that stimulate B cells, T cells, macrophages and mast cells and promote polarization of CD4+ cells into TH2 cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the	idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.
				embodiment of the invention includes a method for stimulating (e.g., increasing) IL-4 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-4 production. A highly preferred indication includes asthma. A highly preferred indication includes allergy. A highly preferred	

			<p>invention) to mediate immunomodulation, stimulate immune cells, modulate immune cell polarization, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-4, and the stimulation of immune cells, such as B cells, T cells, macrophages and mast cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):277-283 (1194); Yssel et al., Res Immunol 144(8):610-616 (1993); Bagley et al., Nat Immunol 1(3):257-261 (2000); and van der Graaff</p>	<p>indication includes rhinitis. Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders").</p>
--	--	--	---	--

			<p>et al., Rheumatology (Oxford) 38(3):214-220 (1999), the contents of each of which are herein incorporated by reference in its entirety.</p> <p>Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art.</p> <p>Human T cells are primary human lymphocytes that mature in the thymus and express a T cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>		<p>Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
	HJBCU04	1199	SEAP in SW480		

251					
252	HJBCY35	1200	SEAP in 3T3L1		
252	HJBCY35	1200	Regulation of viability and proliferation of pancreatic beta cells.	Assays for the regulation of viability and proliferation of cells in vitro are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate viability and proliferation of pancreatic beta cells. For example, the Cell Titer-Glo luminescent cell viability assay measures the number of viable cells in culture based on quantitation of the ATP present which signals the presence of metabolically active cells. Exemplary assays that may be used or routinely modified to test regulation of viability and proliferation of pancreatic beta cells by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Friedrichsen BN,	A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as

				<p>et al., Mol Endocrinol, 15(1):136-48 (2001); Huotari MA, et al., Endocrinology, 139(4):1494-9 (1998); Hugl SR, et al., J Biol Chem 1998 Jul 10;273(28):17771-9 (1998), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an X-ray induced rat transplantable insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Asfari et al. Endocrinology 1992 130:167.</p>	<p>described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
252	HJBCY35	1200	Activation of Skeletal Muscle Cell PI3 Kinase	<p>Kinase assay. Kinase assays, for example an GSK-3 kinase assay, for PI3 kinase signal</p>	<p>A highly preferred embodiment of the invention includes a method for</p>

			<p>transduction that regulate glucose metabolism and cell survival are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit glucose metabolism and cell survival. Exemplary assays for PI3 kinase activity that may be used or routinely modified to test PI3 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Nikoulina et al., Diabetes 49(2):263-271 (2000); and Schreyer et al., Diabetes 48(8):1662-1666 (1999), the contents of each of which are herein incorporated by reference in its entirety. Rat myoblast cells that may be used according to these assays are publicly available (e.g.,</p>	<p>increasing muscle cell survival An alternative highly preferred embodiment of the invention includes a method for decreasing muscle cell survival. A preferred embodiment of the invention includes a method for stimulating muscle cell proliferation. In a specific embodiment, skeletal muscle cell proliferation is stimulated. An alternative highly preferred embodiment of the invention includes a method for inhibiting muscle cell proliferation. In a specific embodiment, skeletal muscle cell proliferation is inhibited. A preferred embodiment of the invention includes a method for stimulating muscle cell differentiation. In a specific embodiment, skeletal muscle cell differentiation is stimulated. An alternative highly preferred embodiment of the invention includes a method for inhibiting muscle cell differentiation. In a specific embodiment, skeletal</p>
--	--	--	---	---

				<p>through the ATCC). Exemplary rat myoblast cells that may be used according to these assays include L6 cells. L6 is an adherent rat myoblast cell line, isolated from primary cultures of rat thigh muscle, that fuses to form multinucleated myotubes and striated fibers after culture in differentiation media.</p> <p>muscle cell differentiation is inhibited. Highly preferred indications include disorders of the musculoskeletal system. Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), endocrine disorders (e.g., as described below under "Endocrine Disorders"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), and infection (e.g., as described below under "Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic</p>
--	--	--	--	--

					<p>nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired</p>
--	--	--	--	--	---

					<p>wound healing, infections (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture).</p> <p>An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p> <p>Additional highly preferred indications are disorders of the musculoskeletal system including myopathies, muscular dystrophy, and/or as described herein.</p> <p>Additional highly preferred indications include: myopathy, atrophy, congestive heart failure, cachexia, myxomas, fibromas, congenital cardiovascular abnormalities, heart disease, cardiac arrest,</p>
--	--	--	--	--	--

					heart valve disease, and vascular disease. Highly preferred indications include neoplasms and cancer, such as, rhabdomyoma, rhabdosarcoma, stomach, esophageal, prostate, and urinary cancer. Preferred indications also include breast, lung, colon, pancreatic, brain, and liver cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, hyperplasia, metaplasia, and/or dysplasia.
253	HJMBI18	1201	IL-2 in Human T-cell 293T	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate serum response factors and modulate the	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described
253	HJMBI18	1201	Activation of transcription through serum response element in immune cells (such as natural killer cells).		

				<p>expression of genes involved in growth and upregulate the function of growth-related genes in many cell types. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line,</p>	<p>below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and</p>
--	--	--	--	---	---

				<p>which is a human natural killer cell line with cytolytic and cytotoxic activity.</p>	<p>cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease,</p>
--	--	--	--	---	--

254	HJMBM38	1202	Endothelial Cell Apoptosis	<p>Caspase Apoptosis. Assays for caspase apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote caspase protease-mediated apoptosis. Induction of apoptosis in endothelial cells supporting the vasculature of tumors is associated with tumor regression due to loss of tumor blood supply. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Lee et al., FEBS</p>	<p>cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p> <p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for</p>
-----	---------	------	----------------------------	--	---

				<p>inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular</p>
			<p>Lett 485(2-3): 122-126 (2000); Nor et al., J Vasc Res 37(3): 209-218 (2000); and Karsan and Harlan, J Atheroscler Thromb 3(2): 75-80 (1996); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through commercial sources). Exemplary endothelial cells that may be used according to these assays include bovine aortic endothelial cells (bAEC), which are an example of endothelial cells which line blood vessels and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	

					<p>disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders").</p> <p>Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization.</p> <p>Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders.</p> <p>Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma,</p>
--	--	--	--	--	--

					<p>hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis,</p>
--	--	--	--	--	--

					<p>lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vascularitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment /prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas,</p>
--	--	--	--	--	---

					heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.
255	HJMBT65	1203	Production of MIP1alpha	MIP-1alpha FMAT. Assays for immunomodulatory proteins produced by activated dendritic cells that upregulate monocyte/macrophage and T cell chemotaxis are well	A highly preferred embodiment of the invention includes a method for stimulating MIP1a production. An alternative highly preferred embodiment of the invention

			<p>known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate chemotaxis, and modulate T cell differentiation. Exemplary assays that test for immunomodulatory proteins evaluate the production of chemokines, such as macrophage inflammatory protein 1 alpha (MIP-1a), and the activation of monocytes/macrophages and T cells. Such assays that may be used or routinely modified to test immunomodulatory and chemotaxis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160</p>	<p>includes a method for inhibiting (e.g., reducing) MIP1a production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma,</p>
--	--	--	--	---

				<p>(2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); Drakes et al., Transp Immunol 8(1):17-29 (2000); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>			<p>arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma, and allergy. Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
255	HJMBT65	1203	CD71 in Human T cells				

256	HJMBW30	1204	SEAP in ATP-3T3-L1	Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et	Highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), and immunodeficiencies (e.g., as described below). An additional highly preferred indication is infection (e.g., AIDS, and/or an infectious disease as described below under "Infectious Disease"). Highly preferred indications include neoplastic diseases (e.g., melanoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred
256	HJMBW30	1204	Activation of transcription through NFKB response element in immune cells (such as T-cells).		

			<p>al., Virus Gnes 15(2):105-117 (1997); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p>	<p>indications include neoplasms and cancers, such as, melanoma, renal cell carcinoma, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, suppression of immune reactions to transplanted</p>
--	--	--	---	---

					organs, asthma and allergy.
256	HJMBW30	1204	SEAP in SW480		
257	HJPAD75	1205	Activation of T-Cell p38 or JNK Signaling Pathway.	<p>Kinase assay. JNK and p38 kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit immune cell (e.g. T-cell) proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc</p>	<p>Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described</p>

			<p>Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.</p>	<p>below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.</p>
HJPAD75	1205	Production of IL-6	IL-6 FMAT. IL-6 is produced	A highly preferred

257				<p>by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins</p>	<p>embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response</p>
-----	--	--	--	---	--

				<p>evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in</p>	<p>and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include</p>
--	--	--	--	--	--

				<p>suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
257	HJPAD75	1205	Glucose Production in H4IIE		
257	HJPAD75	1205	Regulation of transcription through the FAS promoter element in hepatocytes	Assays for the regulation of transcription through the FAS promoter element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure,</p>

				<p>agonists or antagonists of the invention) to activate the FAS promoter element in a reporter construct and to regulate transcription of FAS, a key enzyme for lipogenesis. FAS promoter is regulated by many transcription factors including SREBP. Insulin increases FAS gene transcription in livers of diabetic mice. This stimulation of transcription is also somewhat glucose dependent. Exemplary assays that may be used or routinely modified to test for FAS promoter element activity (in hepatocytes) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Xiong, S., et al., Proc Natl Acad Sci U.S.A., 97(8):3948-53 (2000); Roder, K., et al., Eur J Biochem, 260(3):743-51 (1999); Oskouian B, et al., Biochem J, 317 (Pt 1):257-65 (1996); Berger, et al., Gene 66:1-10 (1988); and, Cullen, B., et al., Methods in Enzymol.</p>	<p>nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and</p>
--	--	--	--	---	---

				<p>216:362–368 (1992), the contents of each of which is herein incorporated by reference in its entirety. Hepatocytes that may be used according to these assays, such as H4IIE cells, are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary hepatocytes that may be used according to these assays include rat liver hepatoma cell line(s) inducible with glucocorticoids, insulin, or cAMP derivatives.</p>	<p>disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
257	HJPAD75	1205	SEAP in HIB/CRE		
257	HJPAD75	1205	Proliferation of pre-adipose cells (such as 3T3-L1 cells)	<p>Assays for the regulation (i.e. increases or decreases) of viability and proliferation of cells in vitro are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate viability and proliferation of pre-adipose cells and cell lines. For</p>	

					example, the CellTiter-Glo [®] Luminescent Cell Viability Assay (Promega Corp., Madison, WI, USA) can be used to measure the number of viable cells in culture based on quantitation of the ATP present which signals the presence of metabolically active cells. 3T3-L1 is a mouse preadipocyte cell line. It is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation. Cells were differentiated to an adipose-like state before being used in the screen. See Green H and Meuth M., Cell 3: 127-133 (1974), which is herein incorporated by reference in its entirety.	
257	HJPAD75	1205	IFN γ in Human T-cell 2B9		Caspase Apoptosis Rescue. Assays for caspase apoptosis rescue are well known in the art and may be used or routinely modified to assess the ability of the polypeptides of the invention (including antibodies and agonists or	A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell
258	HJPCP42	1206	Protection from Endothelial Cell Apoptosis.			

			<p>antagonists of the invention) to inhibit caspase protease-mediated apoptosis.</p> <p>Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis rescue of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Romeo et al., Cardiovasc Res 45(3): 788-794 (2000); Messmer et al., Br J Pharmacol 127(7): 1633-1640 (1999); and J Atheroscler Thromb 3(2): 75-80 (1996); the contents of each of which are herein incorporated by reference in its entirety.</p> <p>Endothelial cells that may be used according to these assays are publicly available (e.g., through commercial sources).</p> <p>Exemplary endothelial cells that may be used according to these assays include bovine aortic endothelial cells (bAEC), which are an example of endothelial cells which line blood vessels and are involved</p>	<p>growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation.</p> <p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells.</p> <p>A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An</p>
--	--	--	--	---

				<p>in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under</p>
--	--	--	--	--	---

					<p>“Cardiovascular Disorders”). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization. Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma,</p>
--	--	--	--	--	---

					<p>haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud"s disease and Reynaud"s phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also</p>
--	--	--	--	--	---

					<p>include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment/prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune</p>
--	--	--	--	--	--

					<p>Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.</p>
258	HJPCP42	1206	<p>Activation or inhibition of transcription through NFKB response element in immune cells (such as basophils).</p>	<p>This reporter assay measures activation or inhibition of the NFKB signaling pathway in Ku812 human basophil cell line. Assays for the activation or inhibition of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including</p>	

				<p>antibodies and agonists or antagonists of the invention) to regulate NFkB transcription factors and modulate expression of immunomodulatory genes. NFkB is important in the pathogenesis of asthma. Exemplary assays for transcription through the NFkB response element that may be used or routinely modified to test NFkB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Marone et al, Int Arch Allergy Immunol 114(3):207-17 (1997), the contents of each of which are herein incorporated by reference in its entirety. Cells were pretreated with SID supernatants or controls for 15-</p>
--	--	--	--	--

260	HKAAH36	1208	Production of IL-6	<p>18 hours, and then 10 ng/mL of TNF was added to stimulate the NFkB reporter. SEAP activity was measured after 48 hours. Basophils that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human basophil cell lines that may be used according to these assays include Ku812, originally established from a patient with chronic myelogenous leukemia. It is an immature prebasophilic cell line that can be induced to differentiate into mature basophils. See, Kishi et al., Leuk Res. 9:381-390 (1985); Blom et al., Eur J Immunol. 22:2025-32 (1992), where the contents of each are herein incorporated by reference in its entirety.</p> <p>IL-6 FMA T. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity).</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a</p>
-----	---------	------	--------------------	--	--

			<p>IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that</p>	<p>method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory</p>
--	--	--	--	---

				<p>may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS,</p>
--	--	--	--	--	--

					<p>granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
260	HKAAH36	1208	Production of MCP-1	<p>MCP-1 FMAT. Assays for immunomodulatory proteins that are produced by a large variety of cells and act to induce chemotaxis and activation of monocytes and T cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, induce chemotaxis, and modulate immune cell activation.</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) MCP-1 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MCP-1 production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Additional highly preferred indications include</p>

			<p>Exemplary assays that test for immunomodulatory proteins evaluate the production of cell surface markers, such as monocyte chemoattractant protein (MCP), and the activation of monocytes and T cells. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using</p>	<p>inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation,</p>
--	--	--	--	---

				<p>techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>diabetes mellitus, endocarditis, meningitis (bacterial and viral), Lyme Disease, asthma, and allergy Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
260	HKAAH36	1208	IgG in Human B cells SAC		
261	HKAAK02	1209	Production of IL-6	<p>IL-6 FMAT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity).</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a</p>

				<p>IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that</p>	<p>method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory</p>
--	--	--	--	--	---

			<p>may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS,</p>
--	--	--	--	--

					<p>granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
261	HKAAK02	1209	Production of MCP-1	<p>MCP-1 FMAT. Assays for immunomodulatory proteins that are produced by a large variety of cells and act to induce chemotaxis and activation of monocytes and T cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, induce chemotaxis, and modulate immune cell activation.</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) MCP-1 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MCP-1 production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Additional highly preferred indications include</p>

				<p>Exemplary assays that test for immunomodulatory proteins evaluate the production of cell surface markers, such as monocyte chemoattractant protein (MCP), and the activation of monocytes and T cells. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using</p>	<p>inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation,</p>
--	--	--	--	--	---

				<p>techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>diabetes mellitus, endocarditis, meningitis (bacterial and viral), Lyme Disease, asthma, and allergy Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
262	HKABI84	1210	Endothelial Cell Apoptosis	<p>Caspase Apoptosis. Assays for caspase apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote caspase</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred</p>

			<p>protease-mediated apoptosis. Induction of apoptosis in endothelial cells supporting the vasculature of tumors is associated with tumor regression due to loss of tumor blood supply. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Lee et al., FEBS Lett 485(2-3): 122-126 (2000); Nor et al., J Vasc Res 37(3): 209-218 (2000); and Karsan and Harlan, J Atheroscler Thromb 3(2): 75-80 (1996); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through commercial sources). Exemplary endothelial cells that may be used according to these assays include bovine</p>	<p>embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment</p>
--	--	--	---	--

				<p>aortic endothelial cells (bAEC), which are an example of endothelial cells which line blood vessels and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders"). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the</p>
--	--	--	--	---	--

					<p>arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization.</p> <p>Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders.</p> <p>Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred</p>
--	--	--	--	--	---

					indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis,
--	--	--	--	--	--

					<p>cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment /prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and</p>
--	--	--	--	--	---

					immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.
262	HKABI84	1210	Activation of transcription through NFAT response in immune cells (such as T-cells).	Assays for the activation of transcription through the Nuclear Factor of Activated T cells (NFAT) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFAT transcription factors and modulate expression of genes involved in immunomodulatory functions. Exemplary assays for transcription through the NFAT response element that may be used or routinely modified to test NFAT-response element activity of	Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. An

				<p>polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Serfling et al., Biochim Biophys Acta 1498(1):1-18 (2000); De Boer et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Fraser et al., Eur J Immunol 29(3):838-844 (1999); and Yeseen et al., J Biol Chem 268(19):14285-14293 (1993), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the JURKAT cell line, which is a suspension culture of leukemia cells that produce IL-2 when stimulated.</p>	<p>additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous</p>
--	--	--	--	---	---

					<p>disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
262	HKABI84	1210	<p>Activation of transcription through NFKB response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides of the invention (including antibodies and</p>	<p>Highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), and immunodeficiencies (e.g., as described below). An additional highly preferred indication is infection (e.g., AIDS, and/or an infectious disease as described below</p>

				<p>agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Gnes 15(2):105-117 (1997); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Exemplary human T cells, such as the MOLT4, that may be used according to these assays are publicly available (e.g., through the ATCC).</p>	<p>under "Infectious Disease"). Highly preferred indications include neoplastic diseases (e.g., melanoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, melanoma, renal cell carcinoma, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel</p>
--	--	--	--	---	---

					<p>disease, sepsis, neutropenia, neutrophilia, psoriasis, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, suppression of immune reactions to transplanted organs, asthma and allergy.</p>
263	HKABZ65	1211	Production of IL-6	<p>IL-6 FMAT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g.,</p>

				<p>polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of</p>	<p>rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon,</p>
--	--	--	--	---	--

			<p>which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
HKABZ65	1211	Activation of	Kinase assay. JNK and p38	A highly preferred

263	Endothelial Cell p38 or JNK Signaling Pathway.	kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999);	embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) endothelial cell activation. An
-----	--	--	---

				<p>the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are endothelial cells which line venous blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) the activation of and/or inactivating endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis,</p>
--	--	--	--	---	---

					<p>cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders").</p> <p>Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization.</p> <p>Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's</p>
--	--	--	--	--	--

					<p>sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's</p>
--	--	--	--	--	--

					phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment
--	--	--	--	--	---

	HKABZ65	1211	Regulation of apoptosis in	Caspase Apoptosis. Assays for caspase apoptosis are well	<p>/prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.</p> <p>A highly preferred indication is diabetes mellitus.</p>
263					

			pancreatic beta cells.	<p>known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote caspase protease-mediated apoptosis. Apoptosis in pancreatic beta is associated with induction and progression of diabetes. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in: Loweth, AC, et al., FEBS Lett, 400(3):285-8 (1997); Saini, KS, et al., Biochem Mol Biol Int, 39(6):1229-36 (1996); Krautheim, A., et al., Br J Pharmacol, 129(4):687-94 (2000); Chandra J, et al., Diabetes, 50 Suppl 1:S44-7 (2001); Suk K, et al., J Immunol, 166(7):4481-9 (2001); Tejedo J, et al., FEBS</p>	<p>An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine</p>
--	--	--	------------------------	---	---

				<p>Lett, 459(2):238-43 (1999); Zhang, S., et al., FEBS Lett, 455(3):315-20 (1999); Lee et al., FEBS Lett 485(2-3): 122-126 (2000); Nor et al., J Vasc Res 37(3): 209-218 (2000); and Karsan and Harlan, J Atheroscler Thromb 3(2): 75-80 (1996); the contents of each of which are herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include RIN-m. RIN-m is a rat adherent pancreatic beta cell insulinoma cell line derived from a radiation induced transplantable rat islet cell tumor. The cells produce and secrete islet polypeptide hormones, and produce insulin, somatostatin, and possibly glucagon. ATTC: #CRL-2057 Chick et al. Proc. Natl. Acad. Sci. 1977 74:628; AF et al.</p>	<p>Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
--	--	--	--	---	--

264	HKACB56	1212	Myoblast cell proliferation	<p>Proc. Natl. Acad. Sci. 1980 77:3519.</p> <p>Assays for muscle cell proliferation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate or inhibit myoblast cell proliferation. Exemplary assays for myoblast cell proliferation that may be used or routinely modified to test activity of polypeptides and antibodies of the invention (including agonists or antagonists of the invention) include, for example, assays disclosed in: Soeta, C., et al. "Possible role for the c-ski gene in the proliferation of myogenic cells in regenerating skeletal muscles of rats" Dev Growth Differ Apr;43(2):155-64 (2001); Ewton DZ, et al., "IGF binding proteins-4, -5 and -6 may play specialized roles during L6 myoblast proliferation and</p>	<p>Highly preferred indications include diabetes, myopathy, muscle cell atrophy, cancers of muscle (such as, rhabdomyoma, and rhabdosarcoma), cardiovascular disorders (such as congestive heart failure, cachexia, myxomas, fibromas, congenital cardiovascular abnormalities, heart disease, cardiac arrest, heart valve disease, vascular disease, and also as described below under "Cardiovascular Disorders"), stimulating myoblast proliferation, and inhibiting myoblast proliferation.</p>
-----	---------	------	-----------------------------	---	---

				differentiation" J Endocrinol Mar;144(3):539-53 (1995); and, Pampusch MS, et al., "Effect of transforming growth factor beta on proliferation of L6 and embryonic porcine myogenic cells" J Cell Physiol Jun;143(3):524-8 (1990); the contents of each of which are herein incorporated by reference in their entirety. Exemplary myoblast cells that may be used according to these assays include the rat myoblast L6 cell line. Rat myoblast L6 cells are an adherent rat myoblast cell line, isolated from primary cultures of rat thigh muscle, that fuse to form multinucleated myotubes and striated fibers after culture in differentiation media.	
264	HKACB56	1212	SEAP in HIB/CRE		
264	HKACB56	1212	CD152 in Human T cells		
264	HKACB56	1212	Production of IL-5	IL-5 FMAT. Assays for immunomodulatory proteins secreted by TH2 cells, mast cells, basophils, and	
				A highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-5	

				<p>eosinophils that stimulate eosinophil function and B cell Ig production and promote polarization of CD4+ cells into TH2 cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, stimulate immune cell function, modulate B cell Ig production, modulate immune cell polarization, and/or mediate humoral or cell-mediated immunity.</p> <p>Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-5, and the stimulation of eosinophil function and B cell Ig production. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the</p>	<p>production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-5 production. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) immunoglobulin production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) immunoglobulin production. A highly preferred indication includes allergy. A highly preferred indication includes asthma. A highly preferred indication includes rhinitis. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or</p>
--	--	--	--	--	--

			<p>invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Ohshima et al., Blood 92(9):3338-3345 (1998); Jung et al., Eur J Immunol 25(8):2413-2416 (1995); Mori et al., J Allergy Clin Immunol 106(1 Pt 2):558-564 (2000); and Koning et al., Cytokine 9(6):427-436 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to</p>	<p>"Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin's disease,</p>
--	--	--	--	--

				immunomodulatory factors.	acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.
264	HKACB56	1212	IFNg in Human T-cell 2B9		
264	HKACB56	1212	Activation of Endothelial Cell p38 or JNK Signaling Pathway.	Kinase assay. JNK and p38 kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase	A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a

			<p>activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are endothelial cells which line venous blood vessels, and are</p>	<p>method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) endothelial cell activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) the activation of and/or inactivating endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A</p>
--	--	--	--	---

				involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.	highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders"). Highly preferred indications include cardiovascular, endothelial and/or angiogenic
--	--	--	--	--	--

					<p>disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization.</p> <p>Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders.</p> <p>Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also</p>
--	--	--	--	--	---

					include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud"s disease and Reynaud"s phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from
--	--	--	--	--	---

					<p>balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment /prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include</p>
--	--	--	--	--	--

264	HKACB56	1212	Production of RANTES in endothelial cells (such as human umbilical vein endothelial cells (HUV/EC))	RANTES FMAT. Assays for immunomodulatory proteins that induce chemotaxis of T cells, monocytes, and eosinophils are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, induce chemotaxis, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for	autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.
-----	---------	------	---	---	--

				<p>immunomodulatory proteins evaluate the production of cytokines, such as RANTES, and the induction of chemotactic responses in immune cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Cocchi et al., Science 270(5243):1811-1815 (1995); and Robinson et al., Clin Exp Immunol 101(3):398-407 (1995), the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary endothelial cells that may be used according to</p>	
--	--	--	--	--	--

264	HKACB56	1212	Production of VCAM in endothelial cells (such as human umbilical vein endothelial cells (HUVVEC))	these assays include human umbilical vein endothelial cells (HUVVEC), which are endothelial cells which line venous blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.	Highly preferred indications include inflammation (acute and chronic), restenosis, atherosclerosis, asthma and allergy. Highly preferred indications include inflammation and inflammatory disorders, immunological disorders, neoplastic disorders (e.g. cancer/tumorigenesis), and cardiovascular disorders (such as described below under "Immune Activity", "Blood-Related Disorders", "Hyperproliferative Disorders" and/or "Cardiovascular Disorders"). Highly preferred indications include neoplasms and cancers such as, for example, leukemia, lymphoma,
				Assays for measuring expression of VCAM are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate VCAM expression. For example, FMAT may be used to measure the upregulation of cell surface VCAM-1 expression in endothelial cells. Endothelial cells are cells that line blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation. Exemplary	

				endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are available from commercial sources. The expression of VCAM (CD106), a membrane-associated protein, can be upregulated by cytokines or other factors, and contributes to the extravasation of lymphocytes, leucocytes and other immune cells from blood vessels; thus VCAM expression plays a role in promoting immune and inflammatory responses.	melanoma, renal cell carcinoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
264	HKACB56	1212	SEAP in SW480		
265	HKACD58	1213	Regulation of transcription via DMEF1 response element in adipocytes and pre-adipocytes	Assays for the regulation of transcription through the DMEF1 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to activate the DMEF1 response element in a	A highly preferred indication is diabetes mellitus. Additional highly preferred indications include complications associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as

				<p>reporter construct (such as that containing the GLUT4 promoter) and to regulate insulin production. The DMEF1 response element is present in the GLUT4 promoter and binds to MEF2 transcription factor and another transcription factor that is required for insulin regulation of Glut4 expression in skeletal muscle. GLUT4 is the primary insulin-responsive glucose transporter in fat and muscle tissue. Exemplary assays that may be used or routinely modified to test for DMEF1 response element activity (in adipocytes and pre-adipocytes) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Thai, M.V., et al., J Biol Chem, 273(23):14285-92 (1998); Mora, S., et al., J Biol Chem, 275(21):16323-8 (2000); Liu, M.L., et al., J Biol Chem, 269(45):28514-21 (1994); "Identification of a 30-base pair regulatory element</p>	<p>described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hypermolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section</p>
--	--	--	--	---	--

				<p>and novel DNA binding protein that regulates the human GLUT4 promoter in transgenic mice", J Biol Chem. 2000 Aug 4;275(31):23666-73; Berger, et al., Gene 66:1-10 (1988); and, Cullen, B., et al., Methods in Enzymol. 216:362-368 (1992), the contents of each of which is herein incorporated by reference in its entirety. Adipocytes and pre-adipocytes that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include the mouse 3T3-L1 cell line which is an adherent mouse preadipocyte cell line. Mouse 3T3-L1 cells are a continuous substrain of 3T3 fibroblasts developed through clonal isolation. These cells undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation culture conditions.</p>	<p>below, especially of the urinary tract and skin). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
--	--	--	--	---	--

265	HKACD58	1213	IL-2 in Human T cells		
265	HKACD58	1213	Activation of transcription through serum response element in immune cells (such as natural killer cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-related genes in many cell types. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications

				<p>85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and cytotoxic activity.</p>	<p>include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia,</p>
--	--	--	--	---	--

					Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
265	HKACD58	1213		Caspase (+camptothecin) in SW480	
265	HKACD58	1213		Caspase (+paclitaxel) in SW480	
266	HKACH44	1214		Activation of Endothelial Cell p38 or JNK Signaling Pathway.	Kinase assay. JNK and p38 kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in
					A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly

			<p>the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be</p>	<p>preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) endothelial cell activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) the</p>
--	--	--	--	--

				<p>used according to these assays are publicly available (e.g., through the ATCC). Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are endothelial cells which line venous blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>activation of and/or inactivating endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular</p>
--	--	--	--	---	---

					<p>disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under “Cardiovascular Disorders”).</p> <p>Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization.</p> <p>Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma,</p>
--	--	--	--	--	---

				<p>hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis,</p>
--	--	--	--	--

					<p>lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment /prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas,</p>
--	--	--	--	--	---

					heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.
267	HKACM93	1215	DT-ATP in DT-ATP-SW480	Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element	Highly preferred indications include asthma, allergy, hypersensitivity reactions, inflammation, and
267	HKACM93	1215	Activation of transcription through GAS response element in		

			immune cells (such as eosinophils).	are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate gene expression (commonly via STAT transcription factors) involved in a wide variety of cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Henttinen et al., J Immunol 155(10):4582-4587 (1995); the contents of each of which are	inflammatory disorders. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting an eosinophil-mediated immune response and, alternatively, suppressing an eosinophil-mediated immune response.
--	--	--	-------------------------------------	--	---

				<p>herein incorporated by reference in its entirety. Moreover, exemplary assays that may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to activate or inhibit activation of immune cells include assays disclosed and/or cited in: Mayumi M., "EoL-1, a human eosinophilic cell line" Leuk Lymphoma; Jun;7(3):243-50 (1992); Bhattacharya S, "Granulocyte macrophage colony-stimulating factor and interleukin-5 activate STAT5 and induce CIS1 mRNA in human peripheral blood eosinophils" Am J Respir Cell Mol Biol; Mar;24(3):312-6 (2001); and, Du J, et al., "Engagement of the CrkL adapter in interleukin-5 signaling in eosinophils" J Biol Chem; Oct 20;275(42):33167-75 (2000); the contents of each of which are herein incorporated by reference in its</p>
--	--	--	--	--

				<p>entirety. Exemplary cells that may be used according to these assays include eosinophils. Eosinophils are a type of immune cell important in the late stage of allergic reactions; they are recruited to tissues and mediate the inflammatory response of late stage allergic reaction. Increases in GAS mediated transcription in eosinophils is typically a result of STAT activation, normally a direct consequence of interleukin or other cytokine receptor stimulation (e.g. IL3, IL5 or GMCSF).</p>	
267	HKACM93	1215	<p>Activation of transcription through GAS response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Gamma Interferon Activation Site (GAS) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT transcription factors and modulate gene expression involved in a wide variety of</p>	<p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate,</p>

				<p>cell functions. Exemplary assays for transcription through the GAS response element that may be used or routinely modified to test GAS-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Matikainen et al., Blood 93(6):1980-1991 (1999); and Hentinen et al., J Immunol 155(10):4582-4587 (1995), the contents of each of which are herein incorporated by reference in its entirety. Exemplary human T cells, such as the MOLT4 cell line, that may be used according to these assays are publicly available (e.g., through the ATCC).</p>	<p>breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis,</p>
--	--	--	--	---	---

					infections associated with chronic granulomatous disease and malignant osteoporosis, and/or an infectious disease as described below under "Infectious Disease"). An additional preferred indication is idiopathic pulmonary fibrosis. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.
267	HKACM93	1215	Activation of transcription through NFKB response element in immune cells (such	Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified	Highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g.,

			as T-cells).	<p>to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Gnes 15(2):105-117 (1997); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Exemplary human T cells,</p>	<p>as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), and immunodeficiencies (e.g., as described below). An additional highly preferred indication is infection (e.g., AIDS, and/or an infectious disease as described below under "Infectious Disease"). Highly preferred indications include neoplastic diseases (e.g., melanoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, melanoma, renal cell carcinoma, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain,</p>
--	--	--	--------------	---	--

				<p>such as the MOL T4, that may be used according to these assays are publicly available (e.g., through the ATCC).</p>	<p>liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, suppression of immune reactions to transplanted organs, asthma and allergy.</p>
268	HKAEL80	1216	<p>Activation of Natural Killer Cell ERK Signaling Pathway.</p>	<p>Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating natural killer cell proliferation. An alternative highly preferred embodiment of the invention includes a</p>

				<p>modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Natural killer cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary natural</p>	<p>method for inhibiting natural killer cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating natural killer cell differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting natural killer cell differentiation. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity") and infections (e.g., as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or</p>
--	--	--	--	---	--

				<p>killer cells that may be used according to these assays include the human natural killer cell lines (for example, NK-YT cells which have cytolytic and cytotoxic activity) or primary NK cells.</p>	<p>"Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include cancers such as, kidney, melanoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, urinary cancer, lymphoma and leukemias. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Other highly preferred indications include, pancytopenia, leukopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), arthritis, asthma,</p>
--	--	--	--	--	---

269	HKAEV06	1217	Regulation of viability and proliferation of pancreatic beta cells.	Assays for the regulation of viability and proliferation of cells in vitro are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate viability and proliferation of pancreatic beta cells. For example, the Cell Titer-Glo luminescent cell viability assay measures the number of viable cells in culture based on quantitation of the ATP present which signals the presence of metabolically active cells. Exemplary assays that may be used or routinely modified to test regulation of viability and proliferation of pancreatic beta cells by polypeptides of the	AIDS, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, immune reactions to transplanted organs and tissues, endocarditis, meningitis, Lyme Disease, and allergies.
				A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g.,	

			<p>invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Ohtani KI, et al., Endocrinology, 139(1):172-8 (1998); Krauthaim A, et al, Exp Clin Endocrinol Diabetes, 107 (1):29-34 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT T15 Cells. HIT T15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-</p>	<p>heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
--	--	--	--	---

					1777 Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.	
269	HKAEV06	1217	DT-ATP in DT-ATP-SW480			
269	HKAEV06	1217	Insulin Inhibition in H4IIE			
269	HKAEV06	1217	Activation of Transcription		Assays for activation of transcription are well-known in the art and may be used and routinely modified to assess ability of polypeptides of the invention to inhibit or activate transcription. An example of such an assay follows: Cells were pretreated with SID supernatants or controls for 15-18 hours. SEAP activity was measured after 48 hours. LS174T is an epithelial colon adenocarcinoma cell line. Its tumorigenicity in nude mice make cell line LS174T a model for studies on the mechanism of synthesis and secretion of specific tumoral markers in colon cancer. See, Patan et al., Circ Res, 89(8):732-39 (2001), the contents of which are	

269	HKAEV06	1217	<p>Activation of transcription through AP1 response element in immune cells (such as T-cells).</p>	<p>herein incorporated by reference in its entirety.</p> <p>Assays for the activation of transcription through the AP1 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for transcription through the AP1 response element that may be used or routinely modified to test AP1-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol</p>	<p>Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under</p>
-----	---------	------	--	---	---

				<p>18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is an IL-2 and IL-4 responsive suspension-culture cell line.</p>	<p>“Hyperproliferative Disorders”). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin’s disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt’s lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.</p>
270	HKAFK41	1218	Production of ICAM-1	Assays for measuring expression of ICAM-1 are	Preferred embodiments of the invention include using

				<p>well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Rolfe BE, et al., Atherosclerosis, 149(1):99-110 (2000); Panettieri RA Jr, et al., J Immunol, 154(5):2358-2365 (1995); and, Grunstein MM, et al., Am J Physiol Lung Cell Mol Physiol, 278(6):L1154-L1163 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include Aortic Smooth Muscle Cells (AOSMC); such as bovine AOSMC.</p>	<p>polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Vascular Disease, Atherosclerosis, Restenosis, Stroke, and Asthma.</p>
--	--	--	--	---	--

270	HKAFK41	1218	Production of IL-6	<p>IL-6 FMT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-</p>
-----	---------	------	--------------------	--	---

			<p>immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are</p>	<p>mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
--	--	--	--	---

				antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.	Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
270	HKAFK41	1218	Proliferation of pre-adipose cells (such as 3T3-L1 cells)	Assays for the regulation (i.e. increases or decreases) of viability and proliferation of cells in vitro are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or	

				antagonists of the invention) to regulate viability and proliferation of pre-adipose cells and cell lines. For example, the CellTiter-Glo [®] Luminescent Cell Viability Assay (Promega Corp., Madison, WI, USA) can be used to measure the number of viable cells in culture based on quantitation of the ATP present which signals the presence of metabolically active cells. 3T3-L1 is a mouse preadipocyte cell line. It is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation. Cells were differentiated to an adipose-like state before being used in the screen. See Green H and Meuth M., Cell 3: 127-133 (1974), which is herein incorporated by reference in its entirety.	
271	HKAF66	1219	Myoblast cell proliferation	Assays for muscle cell proliferation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including	Highly preferred indications include diabetes, myopathy, muscle cell atrophy, cancers of muscle (such as, rhabdomyoma, and rhabdosarcoma),

				<p>antibodies and agonists or antagonists of the invention) to stimulate or inhibit myoblast cell proliferation. Exemplary assays for myoblast cell proliferation that may be used or routinely modified to test activity of polypeptides and antibodies of the invention (including agonists or antagonists of the invention) include, for example, assays disclosed in: Soeta, C., et al. "Possible role for the c-ski gene in the proliferation of myogenic cells in regenerating skeletal muscles of rats" Dev Growth Differ Apr;43(2):155-64 (2001); Ewton DZ, et al., "IGF binding proteins-4, -5 and -6 may play specialized roles during L6 myoblast proliferation and differentiation" J Endocrinol Mar;144(3):539-53 (1995); and, Pampusch MS, et al., "Effect of transforming growth factor beta on proliferation of L6 and embryonic porcine myogenic cells" J Cell Physiol</p>	<p>cardiovascular disorders (such as congestive heart failure, cachexia, myxomas, fibromas, congenital cardiovascular abnormalities, heart disease, cardiac arrest, heart valve disease, vascular disease, and also as described below under "Cardiovascular Disorders"), stimulating myoblast proliferation, and inhibiting myoblast proliferation.</p>
--	--	--	--	---	--

				<p>Jun;143(3):524-8 (1990); the contents of each of which are herein incorporated by reference in their entirety. Exemplary myoblast cells that may be used according to these assays include the rat myoblast L6 cell line. Rat myoblast L6 cells are an adherent rat myoblast cell line, isolated from primary cultures of rat thigh muscle, that fuse to form multinucleated myotubes and striated fibers after culture in differentiation media.</p>	
271	HKAFT66	1219	Insulin Secretion	<p>Assays for measuring secretion of insulin are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin secretion. For example, insulin secretion is measured by FMAT using anti-rat insulin antibodies. Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel</p>	

			<p>proteins/peptides, and dysregulation is a key component in diabetes. Exemplary assays that may be used or routinely modified to test for stimulation of insulin secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Shimizu, H., et al., Endocr J, 47(3):261-9 (2000); Salapatek, A.M., et al., Mol Endocrinol, 13(8):1305-17 (1999); Filipsson, K., et al., Ann N Y Acad Sci, 865:441-4 (1998); Olson, L.K., et al., J Biol Chem, 271(28):16544-52 (1996); and, Miraglia S et al., Journal of Biomolecular Screening, 4:193-204 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that</p>	<p>blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or</p>
--	--	--	---	---

				<p>may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-1777 Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.</p>	<p>complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
271	HKAFT66	1219	<p>Activation of transcription through GATA-3 response element in immune cells (such as mast cells).</p>	<p>This reporter assay measures activation of the GATA-3 signaling pathway in HMC-1 human mast cell line. Activation of GATA-3 in mast cells has been linked to cytokine and chemokine production. Assays for the activation of transcription through the GATA3 response element are well-known in the art and may be used or</p>	<p>Highly preferred indications include allergy, asthma, and rhinitis. Additional preferred indications include infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications also include blood disorders (e.g., as described below under</p>

				<p>routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate GATA3 transcription factors and modulate expression of mast cell genes important for immune response development. Exemplary assays for transcription through the GATA3 response element that may be used or routinely modified to test GATA3-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Flavell et al., Cold Spring Harb Symp Quant Biol 64:563-571 (1999); Rodriguez-Palmero et al., Eur J Immunol 29(12):3914-3924 (1999); Zheng and Flavell, Cell 89(4):587-596 (1997); and</p>	<p>"Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary tract cancers and/or as described below under "Hyperproliferative Disorders"). Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia</p>
--	--	--	--	--	---

				<p>Henderson et al., Mol Cell Biol 14(6):4286-4294 (1994), the contents of each of which are herein incorporated by reference in its entirety. Mast cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human mast cells that may be used according to these assays include the HMC-1 cell line, which is an immature human mast cell line established from the peripheral blood of a patient with mast cell leukemia, and exhibits many characteristics of immature mast cells.</p>	<p>(ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.</p>
271	HKAFT66	1219	<p>Activation of transcription through NFAT response element in immune cells (such as mast cells).</p>	<p>This reporter assay measures activation of the NFAT signaling pathway in HMC-1 human mast cell line. Activation of NFAT in mast cells has been linked to cytokine and chemokine production. Assays for the activation of transcription through the Nuclear Factor of Activated T cells (NFAT) response element are well-</p>	<p>Highly preferred indications include allergy, asthma, and rhinitis. Additional preferred indications include infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications also include blood disorders (e.g., as described below under</p>

			<p>known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFAT transcription factors and modulate expression of genes involved in immunomodulatory functions. Exemplary assays for transcription through the NFAT response element that may be used or routinely modified to test NFAT-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); De Boer et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Ali et al., J Immunol 165(12):7215-7223 (2000);</p>	<p>"Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary tract cancers and/or as described below under "Hyperproliferative Disorders"). Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia</p>
--	--	--	--	---

					Hutchinson and McCloskey, J Biol Chem 270(27):16333-16338 (1995), and Turner et al., J Exp Med 188:527-537 (1998), the contents of each of which are herein incorporated by reference in its entirety. Mast cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human mast cells that may be used according to these assays include the HMC-1 cell line, which is an immature human mast cell line established from the peripheral blood of a patient with mast cell leukemia, and exhibits many characteristics of immature mast cells.	(ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.
271	HKAFT66	1219		IFNg in Human T-cell 2B9		
271	HKAFT66	1219		IL-10 in Human T-cell 2B9		
271	HKAFT66	1219		SEAP in Jurkat/IL4 promoter		
271	HKAFT66	1219		SEAP in Jurkat/IL4 promoter (antiCD3 co-stim)		
	HKDRF34	1220		SEAP in 293/ISRF		

272						
272	HKDBF34	1220	DT-ATP in DT-ATP-SW480			
272	HKDBF34	1220	SEAP in HepG2/Squalene synthetase(stimulated)			
272	HKDBF34	1220	Activation of transcription through NFKB response element in immune cells (such as T-cells).	Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene	Highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), and immunodeficiencies (e.g., as described below). An additional highly preferred indication is infection (e.g., AIDS, and/or an infectious disease as described below under "Infectious Disease"). Highly preferred indications include neoplastic diseases	

				<p>66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Gnes 15(2):105-117 (1997); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Exemplary human T cells, such as the MOLT4, that may be used according to these assays are publicly available (e.g., through the ATCC).</p>	<p>(e.g., melanoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, melanoma, renal cell carcinoma, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, hemophilia, hypercoagulation,</p>
--	--	--	--	---	--

					diabetes mellitus, endocarditis, meningitis, Lyme Disease, suppression of immune reactions to transplanted organs, asthma and allergy.
272	HKDBF34	1220	IL-8 in SW480		
273	HKGAT94	1221	SEAP in ATP-3T3-L1		
273	HKGAT94	1221	Activation of Natural Killer Cell ERK Signaling Pathway.	Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the	A highly preferred embodiment of the invention includes a method for stimulating natural killer cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting natural killer cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating natural killer cell differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting natural killer cell differentiation. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative

				<p>assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Natural killer cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary natural killer cells that may be used according to these assays include the human natural killer cell lines (for example, NK-YT cells which have cytolytic and cytotoxic activity) or primary NK cells.</p>	<p>Disorders”), blood disorders (e.g., as described below under “Immune Activity”, “Cardiovascular Disorders”, and/or “Blood-Related Disorders”), immune disorders (e.g., as described below under “Immune Activity”) and infections (e.g., as described below under “Infectious Disease”). Preferred indications include blood disorders (e.g., as described below under “Immune Activity”, “Blood-Related Disorders”, and/or “Cardiovascular Disorders”). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include cancers such as, kidney, melanoma, prostate,</p>
--	--	--	--	---	--

					breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, urinary cancer, lymphoma and leukemias. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Other highly preferred indications include, pancytopenia, leukopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), arthritis, asthma, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, immune reactions to transplanted organs and tissues, endocarditis, meningitis, Lyme Disease, and allergies.
274	HKGCO27	1222	Production of GM-CSF	GM-CSF FMAT. GM-CSF is expressed by activated T cells, macrophages, endothelial cells, and fibroblasts. GM-CSF regulates differentiation and proliferation of granulocytes-macrophage progenitors and enhances antimicrobial activity	A highly preferred embodiment of the invention includes a method for stimulating the production of GM-CSF. An alternative highly preferred embodiment of the invention includes a method for inhibiting the

				<p>in neutrophils, monocytes and macrophage. Additionally, GM-CSF plays an important role in the differentiation of dendritic cells and monocytes, and increases antigen presentation. GM-CSF is considered to be a proinflammatory cytokine. Assays for immunomodulatory proteins that promote the production of GM-CSF are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and modulate the growth and differentiation of leukocytes. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as GM-CSF, and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention</p>	<p>production of GM-CSF. Highly preferred indications include inflammation and inflammatory disorders. An additional highly preferred indication is infection (e.g., as described below under "Infectious Disease". Highly preferred indications include blood disorders (e.g., neutropenia (and the prevention of neutropenia (e.g., in HIV infected patients), and/or as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications also include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include asthma. Highly preferred indications include neoplastic diseases (e.g., leukemia (e.g., acute</p>
--	--	--	--	--	--

				<p>(including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Ye et al., J Leukoc Biol (58(2):225-233, the contents of each of which are herein incorporated by reference in its entirety.</p> <p>Natural killer cells that may be used according to these assays are publicly available (e.g., through the ATCC) or may be isolated using techniques disclosed herein or otherwise known in the art. Natural killer (NK) cells are large granular lymphocytes that have cytotoxic activity but do bind antigen. NK cells show antibody-independent killing of tumor cells and also recognize antibody bound on target cells, via NK Fc receptors, leading to cell-mediated cytotoxicity.</p>	<p>lymphoblastic leukemia, and acute myelogenous leukemia), lymphoma (e.g., non-Hodgkin's lymphoma and Hodgkin's disease), and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications include: suppression of immune reactions to transplanted organs and tissues (e.g., bone marrow transplant); accelerating myeloid recovery; and mobilizing hematopoietic progenitor cells. Preferred indications include boosting a T cell-mediated immune response, and alternatively,</p>
--	--	--	--	--	---

					suppressing a T cell-mediated immune response. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutrophilia, psoriasis, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and allergy.
275	HKISB57	1223	Activation of JNK Signaling Pathway in immune cells (such as eosinophils).	Kinase assay. JNK kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK kinase activity that may be	Highly preferred indications include asthma, allergy, hypersensitivity reactions, inflammation, and inflammatory disorders. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis

				<p>used or routinely modified to test JNK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Exemplary cells that may be used according to these assays include eosinophils. Eosinophils are important in the late stage of allergic reactions; they are recruited to tissues and mediate the inflammatory response of late stage allergic reaction. Moreover, exemplary assays that may be used or routinely modified to assess the ability</p>	<p>and/or as described below), immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting or inhibiting immune cell proliferation. Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include boosting an eosinophil-mediated immune response, and suppressing an eosinophil-mediated immune response.</p>
--	--	--	--	---	---

				<p>of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate signal transduction, cell proliferation, activation, or apoptosis in eosinophils include assays disclosed and/or cited in: Zhang JP, et al., "Role of caspases in dexamethasone-induced apoptosis and activation of c-Jun NH2-terminal kinase and p38 mitogen-activated protein kinase in human eosinophils" Clin Exp Immunol; Oct;122(1):20-7 (2000); Hebestreit H, et al., "Disruption of fas receptor signaling by nitric oxide in eosinophils" J Exp Med; Feb 2;187(3):415-25 (1998); J Allergy Clin Immunol 1999 Sep;104(3 Pt 1):565-74; and, Sousa AR, et al., "In vivo resistance to corticosteroids in bronchial asthma is associated with enhanced phosphorylation of JUN N-terminal kinase and failure of prednisolone to inhibit JUN N-</p>	
--	--	--	--	--	--

				terminal kinase phosphorylation" J Allergy Clin Immunol; Sep;104(3 Pt 1):565-74 (1999); the contents of each of which are herein incorporated by reference in its entirety.	
275	HKISB57	1223	Regulation of transcription of Malic Enzyme in adipocytes	Assays for the regulation of transcription of Malic Enzyme are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate transcription of Malic Enzyme, a key enzyme in lipogenesis. Malic enzyme is involved in lipogenesis and its expression is stimulated by insulin. ME promoter contains two direct repeat (DR1)- like elements MEp and MEEd identified as putative PPAR response elements. ME promoter may also responds to AP1 and other transcription factors. Exemplary assays that may be used or routinely modified to test for regulation of	A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g.,

				<p>transcription of Malic Enzyme (in adipocytes) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Streeter, R.S., et al., Mol Endocrinol, 12(11):1778-91 (1998); Garcia-Jimenez, C., et al., Mol Endocrinol, 8(10):1361-9 (1994); Barroso, I., et al., J Biol Chem, 274(25):17997-8004 (1999); Ijpenberg, A., et al., J Biol Chem, 272(32):20108-20117 (1997); Berger, et al., Gene 66:1-10 (1988); and, Cullen, B., et al., Methods in Enzymol. 216:362-368 (1992), the contents of each of which is herein incorporated by reference in its entirety. Hepatocytes that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary hepatocytes that may be used according to these assays includes the H4IIE rat</p>	<p>heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
--	--	--	--	--	---

276	HKMLK53	1224	Activation of JNK Signaling Pathway in immune cells (such as eosinophils).	liver hepatoma cell line. Kinase assay. JNK kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK kinase activity that may be used or routinely modified to test JNK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol	Highly preferred indications include asthma, allergy, hypersensitivity reactions, inflammation, and inflammatory disorders. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting or inhibiting immune cell proliferation. Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include boosting an eosinophil-mediated immune response, and suppressing an
-----	---------	------	--	--	--

					<p>Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Exemplary cells that may be used according to these assays include eosinophils. Eosinophils are important in the late stage of allergic reactions; they are recruited to tissues and mediate the inflammatory response of late stage allergic reaction. Moreover, exemplary assays that may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate signal transduction, cell proliferation, activation, or apoptosis in eosinophils include assays disclosed and/or cited in: Zhang JP, et al., "Role of caspases in dexamethasone-induced apoptosis and activation of c-Jun NH2-terminal kinase and p38 mitogen-activated protein kinase in human eosinophils"</p>	eosinophil-mediated immune response.
--	--	--	--	--	---	--------------------------------------

				<p>Clin Exp Immunol; Oct;122(1):20-7 (2000); Hebestreit H, et al., "Disruption of fas receptor signaling by nitric oxide in eosinophils" J Exp Med; Feb 2;187(3):415-25 (1998); J Allergy Clin Immunol 1999 Sep;104(3 Pt 1):565-74; and, Sousa AR, et al., "In vivo resistance to corticosteroids in bronchial asthma is associated with enhanced phosphorylation of JUN N-terminal kinase and failure of prednisolone to inhibit JUN N-terminal kinase phosphorylation" J Allergy Clin Immunol; Sep;104(3 Pt 1):565-74 (1999); the contents of each of which are herein incorporated by reference in its entirety.</p>	
277	HKMLM11	1225	Myoblast cell proliferation	<p>Assays for muscle cell proliferation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to</p> <p>Highly preferred indications include diabetes, myopathy, muscle cell atrophy, cancers of muscle (such as, rhabdomyoma, and rhabdomyosarcoma), cardiovascular disorders (such as congestive heart failure,</p>	

				<p>stimulate or inhibit myoblast cell proliferation. Exemplary assays for myoblast cell proliferation that may be used or routinely modified to test activity of polypeptides and antibodies of the invention (including agonists or antagonists of the invention) include, for example, assays disclosed in: Soeta, C., et al. "Possible role for the c-ski gene in the proliferation of myogenic cells in regenerating skeletal muscles of rats" Dev Growth Differ Apr;43(2):155-64 (2001); Ewton DZ, et al., "IGF binding proteins-4, -5 and -6 may play specialized roles during L6 myoblast proliferation and differentiation" J Endocrinol Mar;144(3):539-53 (1995); and, Pampusch MS, et al., "Effect of transforming growth factor beta on proliferation of L6 and embryonic porcine myogenic cells" J Cell Physiol Jun;143(3):524-8 (1990); the contents of each of which are</p>	<p>cachexia, myxomas, fibromas, congenital cardiovascular abnormalities, heart disease, cardiac arrest, heart valve disease, vascular disease, and also as described below under "Cardiovascular Disorders"), stimulating myoblast proliferation, and inhibiting myoblast proliferation.</p>
--	--	--	--	--	--

				herein incorporated by reference in their entirety. Exemplary myoblast cells that may be used according to these assays include the rat myoblast L6 cell line. Rat myoblast L6 cells are an adherent rat myoblast cell line, isolated from primary cultures of rat thigh muscle, that fuse to form multinucleated myotubes and striated fibers after culture in differentiation media.	
278	HKMLP68	1226	Activation of transcription through serum response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases

				<p>the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>(e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic,</p>
--	--	--	--	---	---

					<p>esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
--	--	--	--	--	--

				<p>used according to these assays are publicly available (e.g., through commercial sources). Exemplary endothelial cells that may be used according to these assays include bovine aortic endothelial cells (bAEC), which are an example of endothelial cells which line blood vessels and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis,</p>
--	--	--	--	---	--

					<p>cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders").</p> <p>Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization.</p> <p>Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's</p>
--	--	--	--	--	--

					<p>sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's</p>
--	--	--	--	--	--

					phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment
--	--	--	--	--	---

279	HKMMD13	1227	Inhibition of squalene synthetase	Reporter Assay: construct contains regulatory and coding	<p>/prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.</p>
-----	---------	------	-----------------------------------	--	--

			gene transcription.	sequence of squalene synthetase, the first specific enzyme in the cholesterol biosynthetic pathway. See Jiang, et al., J. Biol. Chem. 268:12818-12824(1993), the contents of which are herein incorporated by reference in its entirety. Cells were treated with SID supernatants, and SEAP activity was measured after 72 hours. HepG2 is a human hepatocellular carcinoma cell line (ATCC HB-8065). See Knowles et al., Science. 209:497-9 (1980), the contents of which are herein incorporated by reference in its entirety.	
280	HKMND01	1228	TNF α in Human T-cell 293T		
280	HKMND01	1228	Activation of Endothelial Cell p38 or JNK Signaling Pathway.	Kinase assay. JNK and p38 kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or	A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention

				<p>antagonists of the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary endothelial cells that may be used according to</p>	<p>includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) endothelial cell activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) the activation of and/or inactivating endothelial cells. A highly preferred embodiment of the invention includes a method for</p>
--	--	--	--	---	---

				<p>these assays include human umbilical vein endothelial cells (HUVEC), which are endothelial cells which line venous blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or</p>
--	--	--	--	---	---

					<p>as described below under “Cardiovascular Disorders”).</p> <p>Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization.</p> <p>Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders.</p> <p>Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma,</p>
--	--	--	--	--	---

					<p>angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly</p>
--	--	--	--	--	--

					<p>preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment /prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as</p>
--	--	--	--	--	--

					described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.
280	HKMND01	1228	MCP-1 in HUVEC		
280	HKMND01	1228	SEAP in OE-21		
280	HKMND01	1228	SEAP in UMR-106		
281	HL2AC08	1229	IL-2 in Human T-cell 293T		
282	HL2AG57	1230	Production of IL-6	IL-6 FMAT. IL-6 is produced by T cells and has strong effects on B cells. IL-6	A highly preferred embodiment of the invention includes a method for

				<p>participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and</p>	<p>stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune</p>
--	--	--	--	--	---

				<p>the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen</p>	<p>response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia,</p>
--	--	--	--	---	---

				and/or cytokines, initiate and upregulate T cell proliferation and functional activities.	Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
283	HLCND09	1231	DT-ATP in DT-ATP-SW480		
283	HLCND09	1231	SEAP in HIB/CRE		
283	HLCND09	1231	CD152 in Human T cells		
284	HLDBE54	1232	Protection from Endothelial Cell Apoptosis.	Caspase Apoptosis Rescue. Assays for caspase apoptosis rescue are well known in the art and may be used or routinely modified to assess the ability of the polypeptides	A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the

				<p>of the invention (including antibodies and agonists or antagonists of the invention) to inhibit caspase protease-mediated apoptosis. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis rescue of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Romeo et al., Cardiovasc Res 45(3): 788-794 (2000); Messmer et al., Br J Pharmacol 127(7): 1633-1640 (1999); and J Atheroscler Thromb 3(2): 75-80 (1996); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through commercial sources). Exemplary endothelial cells that may be used according to these assays include bovine aortic endothelial cells (bAEC), which are an example</p>	<p>invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention</p>
--	--	--	--	---	--

				<p>of endothelial cells which line blood vessels and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic</p>
--	--	--	--	---	--

					<p>hemodynamic overload, and/or as described below under “Cardiovascular Disorders”).</p> <p>Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization.</p> <p>Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders.</p> <p>Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis,</p>
--	--	--	--	--	--

hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease,					
--	--	--	--	--	--

					<p>and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment /prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood</p>
--	--	--	--	--	---

					disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.
284	HLDBE54	1232	Caspase (+paclitaxel) in SW480		
285	HLDBX13	1233	Production of TNF alpha by dendritic cells	TNFα FMAT. Assays for immunomodulatory proteins produced by activated macrophages, T cells, fibroblasts, smooth muscle, and other cell types that exert a wide variety of inflammatory	A highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) TNF alpha production. An alternative highly preferred embodiment of the invention

				<p>and cytotoxic effects on a variety of cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate inflammation and cytotoxicity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines such as tumor necrosis factor alpha (TNFa), and the induction or inhibition of an inflammatory or cytotoxic response. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160</p>	<p>includes a method for stimulating (e.g., increasing) TNF alpha production. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma,</p>
--	--	--	--	---	---

				<p>(2000); Verhasselt et al., Eur J Immunol 28(11):3886-3890 (1198); Dahlen et al., J Immunol 160(7):3585-3593 (1998); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted</p>
--	--	--	--	--	--

285	HLD BX13	1233	Activation of Endothelial Cell p38 or JNK Signaling Pathway.	<p>Kinase assay. JNK and p38 kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of</p>	<p>organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p> <p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of</p>
-----	----------	------	--	---	---

				<p>the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are endothelial cells which line venous blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) endothelial cell activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) the activation of and/or inactivating endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a</p>
--	--	--	--	---	---

					<p>method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under “Hyperproliferative Disorders”), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under “Cardiovascular Disorders”).</p> <p>Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins</p>
--	--	--	--	--	--

					<p>and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization.</p> <p>Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders.</p> <p>Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign</p>
--	--	--	--	--	---

					<p>dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal</p>
--	--	--	--	--	---

					<p>diseases such as acute renal failure, and osteoporosis.</p> <p>Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment/prevention of endometriosis and related conditions.</p> <p>Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease.</p> <p>Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders").</p> <p>Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as</p>
--	--	--	--	--	--

					described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.
285	HLD BX13	1233	VEGF in SW480		
286	HLD NA86	1234	Activation of Adipocyte ERK Signaling Pathway	<p>Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A highly preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. A highly preferred embodiment of the invention includes a method</p>

				<p>and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to</p>	<p>for stimulating (e.g., increasing) adipocyte activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of (e.g., decreasing) and/or inactivating adipocytes. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"). Highly preferred indications also include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural</p>
--	--	--	--	--	--

				<p>adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease").</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis,</p>
--	--	--	--	---	--

					<p>microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below (particularly of the urinary tract and skin). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems</p>
--	--	--	--	--	---

			through the PEPCK promoter in hepatocytes	<p>PEPCK promoter are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to activate the PEPCK promoter in a reporter construct and regulate liver gluconeogenesis. Exemplary assays for regulation of transcription through the PEPCK promoter that may be used or routinely modified to test for PEPCK promoter activity (in hepatocytes) of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Lochhead et al., Diabetes 49(6):896-903 (2000); and Yeagley et al., J Biol Chem 275(23):17814-17820 (2000),</p>	<p>An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine</p>
--	--	--	---	---	---

				<p>the contents of each of which is herein incorporated by reference in its entirety. Hepatocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary liver hepatoma cells that may be used according to these assays include H4Ile cells, which contain a tyrosine amino transferase that is inducible with glucocorticoids, insulin, or cAMP derivatives.</p>	<p>Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., an infectious diseases or disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include glycogen storage disease (e.g.,</p>
--	--	--	--	--	---

					glycogenoses), hepatitis, gallstones, cirrhosis of the liver, degenerative or necrotic liver disease, alcoholic liver diseases, fibrosis, liver regeneration, metabolic disease, dyslipidemia and cholesterol metabolism, and hepatocarcinomas. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), infection (e.g., an infectious disease and/or disorder as described below under "Infectious Disease"), endocrine disorders (e.g., as described below under "Endocrine Disorders"), and neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"). Additional preferred indications include neoplastic diseases (e.g., as described
--	--	--	--	--	--

					below under “Hyperproliferative Disorders”). Preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, and urinary cancer. A highly preferred indication is liver cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
287	HLDON23	1235	Production of VCAM in endothelial cells (such as human umbilical vein endothelial cells (HUVEC))	Assays for measuring expression of VCAM are well- known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate VCAM expression. For example, FMAT may be used to measure the upregulation of cell surface VCAM-1 expression in endothelial cells. Endothelial cells are cells that line blood	Highly preferred indications include inflammation (acute and chronic), restnosis, atherosclerosis, asthma and allergy. Highly preferred indications include inflammation and inflammatory disorders, immunological disorders, neoplastic disorders (e.g. cancer/tumorigenesis), and cardiovascular disorders (such as described below under “Immune Activity”, “Blood- Related Disorders”,

				<p>vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation. Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are available from commercial sources. The expression of VCAM (CD106), a membrane-associated protein, can be upregulated by cytokines or other factors, and contributes to the extravasation of lymphocytes, leucocytes and other immune cells from blood vessels; thus VCAM expression plays a role in promoting immune and inflammatory responses.</p>	<p>"Hyperproliferative Disorders" and/or "Cardiovascular Disorders"). Highly preferred indications include neoplasms and cancers such as, for example, leukemia, lymphoma, melanoma, renal cell carcinoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
287	HLDON23	1235	Production of ICAM-1	<p>Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and</p>	<p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of</p>

				agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Takacs P, et al, FASEB J, 15(2):279-281 (2001); and, Miyamoto K, et al., Am J Pathol, 156(5):1733-1739 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include microvascular endothelial cells (MVEC).	Inflammation, Vascular Disease, Atherosclerosis, Restenosis, and Stroke
287.	HLDON23	1235	Production of IL-10 and activation of T-cells.	Assays for production of IL-10 and activation of T-cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate or	Highly preferred indications include allergy and asthma. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g.,

				<p>inhibit production of IL-10 and/or activation of T-cells. Exemplary assays that may be used or routinely modified to assess the ability of polypeptides and antibodies of the invention (including agonists or antagonists of the invention) to modulate IL-10 production and/or T-cell proliferation include, for example, assays such as disclosed and/or cited in: Robinson, DS, et al., "Th-2 cytokines in allergic disease" Br Med Bull; 56 (4): 956-968 (2000), and Cohn, et al., "T-helper type 2 cell-directed therapy for asthma" Pharmacology & Therapeutics; 88: 187-196 (2000); the contents of each of which are herein incorporated by reference in their entirety. Exemplary cells that may be used according to these assays include Th2 cells. IL10 secreted from Th2 cells may be measured as a marker of Th2 cell activation. Th2 cells are a class of T cells that secrete</p>	<p>rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response.</p>
--	--	--	--	---	--

				IL4, IL10, IL13, IL5 and IL6. Factors that induce differentiation and activation of Th2 cells play a major role in the initiation and pathogenesis of allergy and asthma. Primary T helper 2 cells are generated via in vitro culture under Th2 polarizing conditions using peripheral blood lymphocytes isolated from cord blood.	
288	HLDOW79	1236	SEAP in HIB/CRE		
288	HLDOW79	1236	Activation of transcription through GATA-3 response element in immune cells (such as mast cells).	<p>This reporter assay measures activation of the GATA-3 signaling pathway in HMC-1 human mast cell line.</p> <p>Activation of GATA-3 in mast cells has been linked to cytokine and chemokine production. Assays for the activation of transcription through the GATA3 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to</p>	<p>Highly preferred indications include allergy, asthma, and rhinitis. Additional preferred indications include infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders.</p> <p>Preferred indications also include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders").</p> <p>Preferred indications include autoimmune diseases (e.g.,</p>

				<p>regulate GATA3 transcription factors and modulate expression of mast cell genes important for immune response development. Exemplary assays for transcription through the GATA3 response element that may be used or routinely modified to test GATA3-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Flavell et al., Cold Spring Harb Symp Quant Biol 64:563-571 (1999); Rodriguez-Palmero et al., Eur J Immunol 29(12):3914-3924 (1999); Zheng and Flavell, Cell 89(4):587-596 (1997); and Henderson et al., Mol Cell Biol 14(6):4286-4294 (1994), the contents of each of which are herein incorporated by reference in its entirety. Mast</p>	<p>rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary tract cancers and/or as described below under "Hyperproliferative Disorders"). Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease,</p>
--	--	--	--	--	--

288	HLDOW79	1236	<p>Activation of transcription through NFAT response element in immune cells (such as mast cells).</p>	<p>cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human mast cells that may be used according to these assays include the HMC-1 cell line, which is an immature human mast cell line established from the peripheral blood of a patient with mast cell leukemia, and exhibits many characteristics of immature mast cells.</p>	<p>sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.</p>
			<p>Activation of transcription through NFAT response element in immune cells (such as mast cells).</p>	<p>This reporter assay measures activation of the NFAT signaling pathway in HMC-1 human mast cell line. Activation of NFAT in mast cells has been linked to cytokine and chemokine production. Assays for the activation of transcription through the Nuclear Factor of Activated T cells (NFAT) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and</p>	<p>Highly preferred indications include allergy, asthma, and rhinitis. Additional preferred indications include infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications also include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g.,</p>

				agonists or antagonists of the invention) to regulate NFAT transcription factors and modulate expression of genes involved in immunomodulatory functions. Exemplary assays for transcription through the NFAT response element that may be used or routinely modified to test NFAT-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); De Boer et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Ali et al., J Immunol 165(12):7215-7223 (2000); Hutchinson and McCloskey, J Biol Chem 270(27):16333-16338 (1995), and Turner et al., J Exp Med 188:527-537 (1998), the contents of each of	rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary tract cancers and/or as described below under "Hyperproliferative Disorders"). Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease,
--	--	--	--	--	---

				<p>which are herein incorporated by reference in its entirety. Mast cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human mast cells that may be used according to these assays include the HMC-1 cell line, which is an immature human mast cell line established from the peripheral blood of a patient with mast cell leukemia, and exhibits many characteristics of immature mast cells.</p>	<p>sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.</p>
288	HLDOW79	1236	<p>Proliferation of pre-adipose cells (such as 3T3-L1 cells)</p>	<p>Assays for the regulation (i.e. increases or decreases) of viability and proliferation of cells in vitro are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate viability and proliferation of pre-adipose cells and cell lines. For example, the CellTiter-Glo[®] Luminescent Cell Viability</p>	

					Assay (Promega Corp., Madison, WI, USA) can be used to measure the number of viable cells in culture based on quantitation of the ATP present which signals the presence of metabolically active cells. 3T3-L1 is a mouse preadipocyte cell line. It is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation. Cells were differentiated to an adipose-like state before being used in the screen. See Green H and Meuth M., Cell 3: 127-133 (1974), which is herein incorporated by reference in its entirety.	
288	HLDOW79	1236	SEAP in Jurkat/IL4 promoter			
288	HLDOW79	1236	SEAP in Jurkat/IL4 promoter (antiCD3 co-stim)			
288	HLDOW79	1236	Activation of transcription through AP1 response element in immune cells (such as T-cells).	Assays for the activation of transcription through the AP1 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention	Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity",	

				<p>(including antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for transcription through the AP1 response element that may be used or routinely modified to test AP1-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays are publicly available (e.g.,</p>	<p>“Cardiovascular Disorders”, and/or “Blood-Related Disorders”), and infection (e.g., an infectious disease as described below under “Infectious Disease”). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under “Hyperproliferative Disorders”). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include</p>
--	--	--	--	---	---

				through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is an IL-2 and IL-4 responsive suspension-culture cell line.	benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.
288	HLDOW79	1236	Activation of transcription through CD28 response element in immune cells (such as T-cells).	Assays for the activation of transcription through the CD28 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate IL-2 expression in T cells.	<p>A highly preferred embodiment of the invention includes a method for stimulating T cell proliferation.</p> <p>An alternative highly preferred embodiment of the invention includes a method for inhibiting T cell proliferation.</p> <p>A highly preferred embodiment of the invention includes a method for</p>

				<p>Exemplary assays for transposition through the CD28 response element that may be used or routinely modified to test CD28-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); McGuire and Iacobelli, J Immunol 159(3):1319-1327 (1997); Parra et al., J Immunol 166(4):2437-2443 (2001); and Butscher et al., J Biol Chem 3(1):552-560 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell</p>	<p>activating T cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating T cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-2 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-2 production. Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Highly preferred</p>
--	--	--	--	---	---

				<p>line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p> <p>indications include neoplastic diseases (e.g., melanoma, renal cell carcinoma, leukemia, lymphoma, and/or as described below under “Hyperproliferative Disorders”). Highly preferred indications include neoplasms and cancers, such as, for example, melanoma (e.g., metastatic melanoma), renal cell carcinoma (e.g., metastatic renal cell carcinoma), leukemia, lymphoma (e.g., T cell lymphoma), and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. A highly preferred indication includes infection (e.g., AIDS, tuberculosis, infections associated with granulomatous disease, and osteoporosis, and/or as described below under “Infectious Disease”). A</p>
--	--	--	--	--

					highly preferred indication is AIDS. Additional highly preferred indications include suppression of immune reactions to transplanted organs and/or tissues, uveitis, psoriasis, and tropical spastic paraparesis. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.
288	HLDOW79	1236	Activation of transcription	Assays for the activation of transcription through the	Highly preferred indications include blood disorders (e.g.,

			through NFAT response element in immune cells (such as T-cells).	<p>Nuclear Factor of Activated T cells (NFAT) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFAT transcription factors and modulate expression of genes involved in immunomodulatory functions. Exemplary assays for transcription through the NFAT response element that may be used or routinely modified to test NFAT-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Serfling et al., Biochim Biophys Acta 1498(1):1-18 (2000); De Boer</p>	<p>as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms</p>
--	--	--	--	---	---

				<p>et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Fraser et al., Eur J Immunol 29(3):838-844 (1999); and Yeseen et al., J Biol Chem 268(19):14285-14293 (1993), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p>	<p>and cancers, such as, for example, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
288	HLDOW79	1236	Activation of transcription	Assays for the activation of transcription through the	Highly preferred indications include inflammation and

			through NFKB response element in immune cells (such as T-cells).	<p>NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely modified to test NFKB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Gnes 15(2):105-117 (1997); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are</p>	<p>inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), and immunodeficiencies (e.g., as described below). An additional highly preferred indication is infection (e.g., AIDS, and/or an infectious disease as described below under "Infectious Disease"). Highly preferred indications include neoplastic diseases (e.g., melanoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, melanoma, renal cell carcinoma, leukemia,</p>
--	--	--	--	--	--

				<p>herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p>	<p>lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, suppression of immune reactions to transplanted organs, asthma and allergy.</p>
289	HLDQC46	1237	Activation of transcription through serum	Assays for the activation of transcription through the Serum Response Element	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g.,</p>

			<p>response element in immune cells (such as T-cells).</p>	<p>(SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used</p>	<p>reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly</p>
--	--	--	--	--	---

				<p>according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous</p>
--	--	--	--	--	--

					<p>disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
290	HLDQR62	1238	<p>Regulation of viability and proliferation of pancreatic beta cells.</p>	<p>Assays for the regulation of viability and proliferation of cells in vitro are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate viability and proliferation of pancreatic beta cells. For example, the Cell Titer-Glo luminescent cell viability assay measures the number of viable cells in culture based on quantitation</p>	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel</p>

				<p>of the ATP present which signals the presence of metabolically active cells. Exemplary assays that may be used or routinely modified to test regulation of viability and proliferation of pancreatic beta cells by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Friedrichsen BN, et al., Mol Endocrinol, 15(1):136-48 (2001); Huotari MA, et al., Endocrinology, 139(4):1494-9 (1998); Hugl SR, et al., J Biol Chem 1998 Jul 10;273(28):17771-9 (1998), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established</p>	<p>blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hypermolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or</p>
--	--	--	--	---	---

290	HLDQR62	1238	Activation of transcription through cAMP response element in immune cells (such as T-cells).	from cells isolated from an X-ray induced rat transplantable insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Asfari et al. Endocrinology 1992 130:167.	Assays for the activation of transcription through the cAMP response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to increase cAMP and regulate CREB transcription factors, and modulate expression of genes involved in a wide variety of cell functions. Exemplary assays for transcription through the cAMP response element that may be used or routinely modified to test cAMP-response element activity of polypeptides of the invention (including antibodies	complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.
				Activation of transcription through cAMP response element in immune cells (such as T-cells).	Assays for the activation of transcription through the cAMP response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to increase cAMP and regulate CREB transcription factors, and modulate expression of genes involved in a wide variety of cell functions. Exemplary assays for transcription through the cAMP response element that may be used or routinely modified to test cAMP-response element activity of polypeptides of the invention (including antibodies	Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional preferred indications include

				<p>and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Genes 15(2):105-117 (1997); and Belkowski et al., J Immunol 161(2):659-665 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTL cell line, which is a suspension culture of IL-2 dependent cytotoxic T cells.</p>	<p>inflammation and inflammatory disorders. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma (e.g., T cell lymphoma, Burkitt's lymphoma, non-Hodgkins lymphoma, Hodgkin's disease), melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, arthritis,</p>
--	--	--	--	---	---

					<p>AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, and asthma and allergy.</p>
291	HLDQU79	1239	Regulation of viability and proliferation of pancreatic beta cells.	<p>Assays for the regulation of viability and proliferation of cells in vitro are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate viability and proliferation of pancreatic beta cells. For example, the Cell Titer-Glo luminescent cell viability assay measures the number of viable cells in culture based on quantitation of the ATP present which signals the presence of metabolically active cells. Exemplary assays that may be</p>	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental</p>

				<p>used or routinely modified to test regulation of viability and proliferation of pancreatic beta cells by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Friedrichsen BN, et al., Mol Endocrinol, 15(1):136-48 (2001); Huotari MA, et al., Endocrinology, 139(4):1494-9 (1998); Hugl SR, et al., J Biol Chem 1998 Jul 10;273(28):17771-9 (1998), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an X-ray induced rat transplantable insulinoma. These cells retain characteristics typical of native</p>	<p>confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively,</p>
--	--	--	--	---	---

				pancreatic beta cells including glucose inducible insulin secretion. References: Asfari et al. Endocrinology 1992 130:167.	weight gain. Additional highly preferred indications are complications associated with insulin resistance.
291	HLDQU79	1239	Activation of transcription through serum response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al.,	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and

				<p>Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
--	--	--	--	--	---

					Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
292	HLDRM43	1240	CD152 in Human T cells		
292	HLDRM43	1240	Activation of Endothelial Cell p38 or JNK Signaling Pathway.	Kinase assay. JNK and p38 kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or	A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the

				<p>routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays</p>	<p>invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) endothelial cell activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) the activation of and/or</p>
--	--	--	--	--	--

				<p>are publicly available (e.g., through the ATCC). Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are endothelial cells which line venous blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>inactivating endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy,</p>
--	--	--	--	--	---

					intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders"). Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization. Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and
--	--	--	--	--	---

					<p>cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and</p>
--	--	--	--	--	--

					<p>lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment /prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest,</p>
--	--	--	--	--	--

292	HDRM43	1240	Production of RANTES in endothelial cells (such as human umbilical vein endothelial cells (HUVEC))	RANTES FMAT. Assays for immunomodulatory proteins that induce chemotaxis of T cells, monocytes, and eosinophils are well known in the art and may be used or routinely modified to assess	heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.
-----	--------	------	--	---	--

				<p>the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, induce chemotaxis, and/or mediate humoral or cell-mediated immunity.</p> <p>Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as RANTES, and the induction of chemotactic responses in immune cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Cocchi et al., Science 270(5243):1811-1815 (1995); and Robinson et al., Clin Exp Immunol 101(3):398-407</p>
--	--	--	--	---

292	HLD43	1240	<p>Activation of transcription through serum response element in immune cells (such as natural killer cells).</p>	<p>(1995), the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are endothelial cells which line venous blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p> <p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate serum response factors and modulate the</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described</p>
-----	-------	------	---	---	---

				<p>expression of genes involved in growth and upregulate the function of growth-related genes in many cell types. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line,</p>	<p>below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and</p>
--	--	--	--	---	---

				<p>which is a human natural killer cell line with cytolytic and cytotoxic activity.</p>	<p>cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease,</p>
--	--	--	--	---	--

					cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
292	HILDRM43	1240	Activation of transcription through CD28 response element in immune cells (such as T-cells).	Assays for the activation of transcription through the CD28 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate IL-2 expression in T cells. Exemplary assays for transcription through the CD28 response element that may be used or routinely modified to test CD28-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA	<p>A highly preferred embodiment of the invention includes a method for stimulating T cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting T cell proliferation. A highly preferred embodiment of the invention includes a method for activating T cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating T cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-2 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g.,</p>

				<p>85:6342-6346 (1988); McGuire and Iacobelli, J Immunol 159(3):1319-1327 (1997); Parra et al., J Immunol 166(4):2437-2443 (2001); and Butscher et al., J Biol Chem 3(1):552-560 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p>	<p>reducing) IL-2 production. Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Highly preferred indications include neoplastic diseases (e.g., melanoma, renal cell carcinoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, melanoma (e.g., metastatic melanoma), renal cell carcinoma (e.g., metastatic renal cell carcinoma), leukemia, lymphoma (e.g., T</p>
--	--	--	--	--	--

					<p>cell lymphoma), and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. A highly preferred indication includes infection (e.g., AIDS, tuberculosis, infections associated with granulomatous disease, and osteoporosis, and/or as described below under "Infectious Disease"). A highly preferred indication is AIDS. Additional highly preferred indications include suppression of immune reactions to transplanted organs and/or tissues, uveitis, psoriasis, and tropical spastic paraparesis. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders").</p>
--	--	--	--	--	--

					<p>Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
293	HLD RP33	1241	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or</p>	<p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"),</p>

				<p>routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid</p>
--	--	--	--	--	---

					<p>tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious</p>
--	--	--	--	--	---

294	HLJHFP03	1242	Activation of T-Cell p38 or JNK Signaling Pathway.	<p>Kinase assay. JNK and p38 kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit immune cell (e.g. T-cell) proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang</p>	<p>disease as described below under "Infectious Disease"). Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under</p>
-----	----------	------	--	---	---

				and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTL cell line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.	"Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.
294	HLHFP03	1242	SEAP in HIB/CRE		
	HLHFP03	1242	VEGF in HT1080		

294	HLHFP03	1242	Production of VCAM in endothelial cells (such as human umbilical vein endothelial cells (HUVEC))	Assays for measuring expression of VCAM are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate VCAM expression. For example, FMAT may be used to measure the upregulation of cell surface VCAM-1 expression in endothelial cells. Endothelial cells are cells that line blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation. Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are available from commercial sources. The expression of VCAM (CD106), a membrane-associated protein, can be	Highly preferred indications include inflammation (acute and chronic), restenosis, atherosclerosis, asthma and allergy. Highly preferred indications include inflammation and inflammatory disorders, immunological disorders, neoplastic disorders (e.g. cancer/tumorigenesis), and cardiovascular disorders (such as described below under "Immune Activity", "Blood-Related Disorders", "Hyperproliferative Disorders" and/or "Cardiovascular Disorders"). Highly preferred indications include neoplasms and cancers such as, for example, leukemia, lymphoma, melanoma, renal cell carcinoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for
-----	---------	------	--	---	---

					upregulated by cytokines or other factors, and contributes to the extravasation of lymphocytes, leucocytes and other immune cells from blood vessels; thus VCAM expression plays a role in promoting immune and inflammatory responses.	example, hyperplasia, metaplasia, and/or dysplasia.
294	HLHFP03	1242	SEAP in OE-21			
295	HLHFR58	1243	Production of TNF alpha by dendritic cells		TNFα F-MAT. Assays for immunomodulatory proteins produced by activated macrophages, T cells, fibroblasts, smooth muscle, and other cell types that exert a wide variety of inflammatory and cytotoxic effects on a variety of cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate inflammation and cytotoxicity. Exemplary assays that test for immunomodulatory proteins	A highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus,

				<p>evaluate the production of cytokines such as tumor necrosis factor alpha (TNFa), and the induction or inhibition of an inflammatory or cytotoxic response. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Verhasselt et al., Eur J Immunol 28(11):3886-3890 (1998); Dahlen et al., J Immunol 160(7):3585-3593 (1998); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these</p>	<p>Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include</p>
--	--	--	--	---	---

				assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.	benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
295	HLHFR58	1243	RANTES in Human T cells		
	HLIBD68	1244	Production of IL-6	IL-6 FMAT. IL-6 is produced	A highly preferred

296				<p>by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins</p>	<p>embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response</p>
-----	--	--	--	---	--

				<p>evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in</p>	<p>and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include</p>
--	--	--	--	--	--

				<p>suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
296	HLIBD68	1244	<p>Production of MIP1alpha</p>	<p>MIP-1alpha FMA T. Assays for immunomodulatory proteins produced by activated dendritic cells that upregulate monocyte/macrophage and T cell chemotaxis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating MIP1a production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MIP1a production. A highly preferred indication is</p>

				<p>(including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate chemotaxis, and modulate T cell differentiation. Exemplary assays that test for immunomodulatory proteins evaluate the production of chemokines, such as macrophage inflammatory protein 1 alpha (MIP-1a), and the activation of monocytes/macrophages and T cells. Such assays that may be used or routinely modified to test immunomodulatory and chemotaxis activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); Drakes et al., Transp Immunol 8(1):17-</p>	<p>infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis,</p>
--	--	--	--	---	--

				<p>29 (2000); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety.</p> <p>Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art.</p> <p>Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma, and allergy.</p> <p>Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
296	HLIBD68	1244	<p>Production of TNF alpha by dendritic cells</p>	<p>TNFα FMAT. Assays for immunomodulatory proteins produced by activated macrophages, T cells, fibroblasts, smooth muscle, and other cell types that exert a</p> <p>A highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) TNF alpha production. An alternative highly preferred</p>	

				<p>wide variety of inflammatory and cytotoxic effects on a variety of cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate inflammation and cytotoxicity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines such as tumor necrosis factor alpha (TNFa), and the induction or inhibition of an inflammatory or cytotoxic response. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical</p>	<p>embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases</p>
--	--	--	--	--	--

				<p>approach" Chapter 6:138-160 (2000); Verhasselt et al., Eur J Immunol 28(11):3886-3890 (1998); Dahlen et al., J Immunol 160(7):3585-3593 (1998); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>(e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune</p>
--	--	--	--	--	---

					reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
296	HLIBD68	1244	Stimulation of insulin secretion from pancreatic beta cells.	Assays for measuring secretion of insulin are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin secretion. For example, insulin secretion is measured by FMAT using anti-rat insulin antibodies. Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain proteins/peptides, and dysregulation is a key component in diabetes. Exemplary assays that may be	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental</p>

				<p>used or routinely modified to test for stimulation of insulin secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Ahren, B., et al., Am J Physiol, 277(4 Pt 2):R959-66 (1999); Li, M., et al., Endocrinology, 138(9):3735-40 (1997); Kim, K.H., et al., FEBS Lett, 377(2):237-9 (1995); and, Miraglia S et. al., Journal of Biomolecular Screening, 4:193-204 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an X-ray induced</p>	<p>confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively,</p>
--	--	--	--	---	---

297	HLICQ90	1245	<p>Activation of transcription through serum response element in immune cells (such as T-cells).</p>	<p>rat transplantable insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Asfari et al. Endocrinology 1992 130:167.</p> <p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10</p>	<p>weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p> <p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies</p>
-----	---------	------	--	---	---

				<p>(1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>(e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic</p>
--	--	--	--	---	--

					conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
297	HLICQ90	1245	Production of TNF alpha by dendritic cells	TNFa FMAT. Assays for immunomodulatory proteins produced by activated macrophages, T cells, fibroblasts, smooth muscle,	A highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) TNF alpha production. An

				<p>and other cell types that exert a wide variety of inflammatory and cytotoxic effects on a variety of cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate inflammation and cytotoxicity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines such as tumor necrosis factor alpha (TNFa), and the induction or inhibition of an inflammatory or cytotoxic response. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al.,</p>	<p>alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications</p>
--	--	--	--	--	---

				<p>"Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Verhasselt et al., Eur J Immunol 28(11):3886-3890 (1998); Dahlen et al., J Immunol 160(7):3585-3593 (1998); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis,</p>
--	--	--	--	--	---

297	HLICQ90	1245	Stimulation of Calcium Flux in pancreatic beta cells.	Assays for measuring calcium flux are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mobilize calcium. For example, the FLPR assay may be used to measure influx of calcium. Cells normally have very low concentrations of cytosolic calcium compared to much higher extracellular calcium. Extracellular factors can cause an influx of calcium, leading to activation of calcium responsive signaling pathways	suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
				A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel	

				<p>and alterations in cell functions. Exemplary assays that may be used or routinely modified to measure calcium flux by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Satin LS, et al., Endocrinology, 136(10):4589-601 (1995); Mogami H, et al., Endocrinology, 136(7):2960-6 (1995); Richardson SB, et al., Biochem J, 288 (Pt 3):847-51 (1992); and, Meats, JE, et al., Cell Calcium 1989 Nov-Dec;10(8):535-41 (1989), the contents of each of which is herein incorporated by reference in its entirety.</p> <p>Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT T15 Cells. HIT T15 are an adherent epithelial cell line established from Syrian hamster islet cells</p>	<p>blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include</p>
--	--	--	--	---	---

297	HILICQ90	1245	<p>transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-1777 Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.</p> <p>Assays for measuring secretion of insulin are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin secretion. For example, insulin secretion is measured by FMAT using anti-rat insulin antibodies. Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain proteins/peptides, and</p>	<p>weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke,</p>
-----	----------	------	---	---

			<p>disregulation is a key component in diabetes. Exemplary assays that may be used or routinely modified to test for stimulation of insulin secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Ahren, B., et al., Am J Physiol, 277(4 Pt 2):R959-66 (1999); Li, M., et al., Endocrinology, 138(9):3735-40 (1997); Kim, K.H., et al., FEBS Lett, 377(2):237-9 (1995); and, Miraglia S et. al., Journal of Biomolecular Screening, 4:193-204 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1</p>	<p>impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with</p>
--	--	--	--	--

298	HLMBO76	1246	Activation of transcription through serum response element in immune cells (such as natural killer cells).	cells are a semi-adherent cell line established from cells isolated from an X-ray induced rat transplantable insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Asfari et al. Endocrinology 1992 130:167.	obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.
			Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-related genes in many cell types. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus,	

				<p>of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and cytotoxic activity.</p>	<p>Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammatory and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other</p>
--	--	--	--	--	--

					<p>preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
299	HLQBE09	1247	Production of IL-8 by immune cells	Assay that measures the production of the chemokine	Highly preferred indications include eosinophilia, asthma,

			(such as the human EOL-1 eosinophil cells)	<p>interleukin-8 (IL-8) from immune cells (such as the EOL-1 human eosinophil cell line) are well known in the art (for example, measurement of IL-8 production by FMAT) and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit. Eosinophils are a type of immune cell important in allergic responses; they are recruited to tissues and mediate the inflammatory response of late stage allergic reaction. IL8 is a strong immunomodulator and may have a potential proinflammatory role in immunological diseases and disorders (such as allergy and asthma).</p>	<p>allergy, hypersensitivity reactions, inflammation, and inflammatory disorders. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting or inhibiting immune cell proliferation. Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include boosting an eosinophil-mediated immune response, and suppressing an eosinophil-mediated immune response.</p>	
	HLQBE09	1247	SEAP in HIB/CRE			

299						
299	HLQBE09	1247	CD71 in Human T cells			
299	HLQBE09	1247	IL-10 in Human T-cell 293T			
299	HLQBE09	1247	TNFa in Human T-cell 2B9			
300	HLQDR48	1248	Activation of Adipocyte ERK Signaling Pathway	Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-	A highly preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A highly preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) adipocyte activation. An alternative highly preferred embodiment	

				<p>1110 (1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>of the invention includes a method for inhibiting the activation of (e.g., decreasing) and/or inactivating adipocytes. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"). Highly preferred indications also include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as</p>
--	--	--	--	--	--

					<p>described below under "Infectious Disease").</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the</p>
--	--	--	--	--	--

					<p>"Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below (particularly of the urinary tract and skin). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred</p>
--	--	--	--	--	---

					indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Preferred indications include neoplasms and cancer, such as, lymphoma, leukemia and breast, colon, and kidney cancer. Additional preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Highly preferred indications include lipomas and liposarcomas. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
300	HLQDR48	1248	Production of MCP-1	MCP-1 FMAT. Assays for immunomodulatory proteins that are produced by a large variety of cells and act to induce chemotaxis and activation of monocytes and T	A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) MCP-1 production. An alternative highly preferred

			<p>cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, induce chemotaxis, and modulate immune cell activation. Exemplary assays that test for immunomodulatory proteins evaluate the production of cell surface markers, such as monocyte chemoattractant protein (MCP), and the activation of monocytes and T cells. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Sathaporn and</p>	<p>embodiment of the invention includes a method for inhibiting (e.g., reducing) MCP-1 production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL),</p>
--	--	--	--	--

				<p>Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety.</p> <p>Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art.</p> <p>Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis (bacterial and viral), Lyme Disease, asthma, and allergy Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia,</p>
--	--	--	--	--	--

300	HLQDR48	1248	Production of TNF alpha by dendritic cells	<p>TNFα FMAT. Assays for immunomodulatory proteins produced by activated macrophages, T cells, fibroblasts, smooth muscle, and other cell types that exert a wide variety of inflammatory and cytotoxic effects on a variety of cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate inflammation and cytotoxicity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines such as tumor necrosis factor alpha (TNFα), and the induction or inhibition of an inflammatory or cytotoxic response. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention</p>	<p>metaplasia, and/or dysplasia.</p> <p>A highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and</p>
-----	---------	------	--	---	---

				<p>(including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Verhasselt et al., Eur J Immunol 28(11):3886-3890 (1998); Dahlen et al., J Immunol 160(7):3585-3593 (1998); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL),</p>
--	--	--	--	---	---

					<p>plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
300	HLQDR48	1248	IL-8 in Normal Human Bronchial Epitheliae	<p>Assays for the activation of transcription through the API response element are known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and other cell</p>	<p>Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection</p>
301	HLTAU74	1249	<p>Activation of transcription through API response element in immune cells (such as T-cells).</p>		

				<p>functions. Exemplary assays for transcription through the AP1 response element that may be used or routinely modified to test AP1-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these</p>	<p>(e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for</p>
--	--	--	--	--	--

				assays include the CTLL cell line, which is an IL-2 dependent suspension-culture cell line with cytotoxic activity.	example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.
301	HLTAU74	1249	Activation of Natural Killer Cell ERK Signaling Pathway.	Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation.	A highly preferred embodiment of the invention includes a method for stimulating natural killer cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting natural killer cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating natural killer cell differentiation. An alternative highly preferred

				<p>Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Natural killer cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary natural killer cells that may be used according to these assays include the human natural killer cell lines (for example, NK-YT cells which have cytolytic and cytotoxic activity) or primary NK cells.</p>	<p>embodiment of the invention includes a method for inhibiting natural killer cell differentiation. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity") and infections (e.g., as described below under "Infectious Disease"). Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and</p>
--	--	--	--	--	--

					<p>immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders.</p> <p>Highly preferred indications also include cancers such as, kidney, melanoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, urinary cancer, lymphoma and leukemias.</p> <p>Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p> <p>Other highly preferred indications include, pancytopenia, leukopenia, leukemias, Hodgkin's disease, acute lymphocytic anemia (ALL), arthritis, asthma, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, immune reactions to transplanted organs and tissues, endocarditis, meningitis, Lyme Disease, and allergies.</p>
--	--	--	--	--	---

302	HLTDV50	1250	IL-10 in Human T-cell 2B9		Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Takacs P, et al, FASEB J, 15(2):279-281 (2001); and, Miyamoto K, et al., Am J Pathol, 156(5):1733-1739 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include microvascular endothelial cells (MVEC).	Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Inflammation, Vascular Disease, Atherosclerosis, Restenosis, and Stroke
302	HLTDV50	1250	Production of ICAM-1			

303	HL TEI25	1251	<p>Activation of transcription through GATA-3 response element in immune cells (such as mast cells).</p>	<p>This reporter assay measures activation of the GATA-3 signaling pathway in HMC-1 human mast cell line.</p> <p>Activation of GATA-3 in mast cells has been linked to cytokine and chemokine production. Assays for the activation of transcription through the GATA3 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate GATA3 transcription factors and modulate expression of mast cell genes important for immune response development. Exemplary assays for transcription through the GATA3 response element that may be used or routinely modified to test GATA3-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays</p>	<p>Highly preferred indications include allergy, asthma, and rhinitis. Additional preferred indications include infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders.</p> <p>Preferred indications also include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders").</p> <p>Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary tract cancers and/or as described below under</p>
-----	----------	------	--	---	---

				disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Flavell et al., Cold Spring Harb Symp Quant Biol 64:563-571 (1999); Rodriguez-Palmero et al., Eur J Immunol 29(12):3914-3924 (1999); Zheng and Flavell, Cell 89(4):587-596 (1997); and Henderson et al., Mol Cell Biol 14(6):4286-4294 (1994), the contents of each of which are herein incorporated by reference in its entirety. Mast cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human mast cells that may be used according to these assays include the HMC-1 cell line, which is an immature human mast cell line established from the peripheral blood of a patient with mast cell leukemia, and exhibits many characteristics of immature mast cells.	“Hyperproliferative Disorders”). Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin’s disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt’s lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.
--	--	--	--	--	--

303	HLTEI25	1251	<p>Activation of transcription through NFAT response element in immune cells (such as mast cells).</p>	<p>This reporter assay measures activation of the NFAT signaling pathway in HMC-1 human mast cell line. Activation of NFAT in mast cells has been linked to cytokine and chemokine production. Assays for the activation of transcription through the Nuclear Factor of Activated T cells (NFAT) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFAT transcription factors and modulate expression of genes involved in immunomodulatory functions. Exemplary assays for transcription through the NFAT response element that may be used or routinely modified to test NFAT-response element activity of polypeptides of the invention (including antibodies and</p>	<p>Highly preferred indications include allergy, asthma, and rhinitis. Additional preferred indications include infection (e.g., an infectious disease as described below under "Infectious Disease"), and inflammation and inflammatory disorders. Preferred indications also include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary tract cancers and/or as described below under</p>
-----	---------	------	--	--	---

				<p>agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); De Boer et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Ali et al., J Immunol 165(12):7215-7223 (2000); Hutchinson and McCloskey, J Biol Chem 270(27):16333-16338 (1995), and Turner et al., J Exp Med 188:527-537 (1998), the contents of each of which are herein incorporated by reference in its entirety. Mast cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human mast cells that may be used according to these assays include the HMC-1 cell line, which is an immature human mast cell line established from the peripheral blood of a patient with mast cell leukemia, and exhibits</p>	<p>“Hyperproliferative Disorders”). Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, leukemias, Hodgkin’s disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt’s lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease.</p>
--	--	--	--	--	---

					many characteristics of immature mast cells.	
303	HLTEI25	1251	IL-6 in HUVEC			
304	HLTEJ06	1252	Activation of transcription through serum response element in immune cells (such as T-cells).		Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated

				<p>85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include</p>
--	--	--	--	---	---

305	HLTFA64	1253	Production of IFNgamma using Natural Killer cells	IFNgamma FMAT. IFNγ plays a central role in the immune system and is considered to be a proinflammatory cytokine. IFNγ promotes TH1 and inhibits TH2; promotes IgG2a and inhibits IgE; induces macrophage activation; and	anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). A highly preferred embodiment of the invention includes a method for stimulating the production of IFNγ. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of IFNγ. Highly preferred
-----	---------	------	---	---	---

				<p>increases MHC expression. Assays for immunomodulatory proteins produced by T cells and NK cells that regulate a variety of inflammatory activities and inhibit TH2 helper cell functions are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon gamma (IFNg), and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and</p>	<p>indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", "Hyperproliferative Disorders" (e.g. cancer/tumorigenesis) and/or "Cardiovascular Disorders"), and infection (e.g., viral infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiency (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response, boosting antibody-dependent immune responses, suppressing antibody-dependent immune responses, boosting innate</p>
--	--	--	--	---	---

				<p>agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated by reference in its entirety. Natural Killer (NK) cells that may be used according to these assays are publicly available (e.g., through the ATCC) or may be isolated using techniques disclosed herein or otherwise known in the art. Natural killer (NK) cells are large granular lymphocytes that have cytotoxic activity but do bind antigen. NK cells show antibody-independent killing of tumor cells and also</p>	<p>immunity and immune responses, and suppressing innate immunity and immune responses. Additional highly preferred indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include</p>
--	--	--	--	--	--

				recognize antibody bound on target cells, via NK Fc receptors, leading to cell-mediated cytotoxicity.	anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.
306	HLTHG37	1254	Glucose Production in H4IIE		
306	HLTHG37	1254	Activation of transcription through STAT6 response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Signal Transducers and Activators of Transcription (STAT6) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate STAT6	A highly preferred indication is allergy. Another highly preferred indication is asthma. Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-

			<p>transcription factors and modulate the expression of multiple genes. Exemplary assays for transcription through the STAT6 response element that may be used or routinely modified to test STAT6 response element activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Georas et al., Blood 92(12):4529-4538 (1998); Moffatt et al., Transplantation 69(7):1521-1523 (2000); Curiel et al., Eur J Immunol 27(8):1982-1987 (1997); and Masuda et al., J Biol Chem 275(38):29331-29337 (2000), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays</p>	<p>Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia,</p>
--	--	--	---	---

				are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.	leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
307	HLWAA17	1255	Regulation of transcription of Malic Enzyme in adipocytes	Assays for the regulation of transcription of Malic Enzyme are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate transcription of Malic Enzyme, a key enzyme in lipogenesis.	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal</p>

				<p>Malic enzyme is involved in lipogenesis and its expression is stimulated by insulin. ME promoter contains two direct repeat (DR1)-like elements MEp and MEEd identified as putative PPAR response elements. ME promoter may also respond to AP1 and other transcription factors. Exemplary assays that may be used or routinely modified to test for regulation of transcription of Malic Enzyme (in adipocytes) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Streeter, R.S., et al., Mol Endocrinol, 12(11):1778-91 (1998); Garcia-Jimenez, C., et al., Mol Endocrinol, 8(10):1361-9 (1994); Barroso, I., et al., J Biol Chem, 274(25):17997-8004 (1999); Ijpenberg, A., et al., J Biol Chem, 272(32):20108-20117 (1997); Berger, et al., Gene 66:1-10 (1988); and, Cullen, B., et al.,</p>	<p>Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the</p>
--	--	--	--	--	---

307	HLWAA17	1255		<p>Methods in Enzymol. 216:362-368 (1992), the contents of each of which is herein incorporated by reference in its entirety. Hepatocytes that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary hepatocytes that may be used according to these assays includes the H4IIE rat liver hepatoma cell line.</p> <p>Assays for measuring expression of ICAM-1 are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate ICAM-1 expression. Exemplary assays that may be used or routinely modified to measure ICAM-1 expression include assays disclosed in: Takacs P, et al, FASEB J, 15(2):279-281 (2001); and, Miyamoto K, et al., Am J Pathol, 156(5):1733-</p>	<p>urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p> <p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Inflammation, Vascular Disease, Atherosclerosis, Restenosis, and Stroke</p>
-----	---------	------	--	---	---

308	HLWAA88	1256	Activation of Endothelial Cell p38 or JNK Signaling Pathway.	<p>1739 (2000), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include microvascular endothelial cells (MVEC).</p> <p>Kinase assay. JNK and p38 kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention</p>
-----	---------	------	--	--	---

				<p>invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are endothelial cells which line venous blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone,</p>	<p>includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) endothelial cell activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) the activation of and/or inactivating endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative</p>
--	--	--	--	---	--

					<p>highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under "Cardiovascular Disorders").</p> <p>Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels</p>
--	--	--	--	--	--

					<p>themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization.</p> <p>Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders.</p> <p>Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and</p>
--	--	--	--	--	--

					<p>urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud"s disease and Reynaud"s phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury,</p>
--	--	--	--	--	---

					<p>rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment/prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described</p>
--	--	--	--	--	---

308	HLWAA88	1256	Production of RANTES in endothelial cells (such as human umbilical vein endothelial cells (HUVEC))	RANTES FMAT. Assays for immunomodulatory proteins that induce chemotaxis of T cells, monocytes, and eosinophils are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, induce chemotaxis, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as RANTES, and the induction of	below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.
-----	---------	------	--	--	--

				<p>chemotactic responses in immune cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Cocchi et al., Science 270(5243):1811-1815 (1995); and Robinson et al., Clin Exp Immunol 101(3):398-407 (1995), the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUVEC), which are endothelial cells which line</p>	
--	--	--	--	---	--

309	HLW/AD77	1257	Proliferation of pre-adipose cells (such as 3T3-L1 cells)	<p>venous blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p> <p>Assays for the regulation (i.e. increases or decreases) of viability and proliferation of cells in vitro are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate viability and proliferation of pre-adipose cells and cell lines. For example, the CellTiter-Glo[®] Luminescent Cell Viability Assay (Promega Corp., Madison, WI, USA) can be used to measure the number of viable cells in culture based on quantitation of the ATP present which signals the presence of metabolically active cells. 3T3-L1 is a mouse preadipocyte cell line. It is a continuous substrain of</p>	
-----	----------	------	---	---	--

309	HLWAD77	1257	<p>Activation of transcription through the EGR (Early Growth Response) element in immune cells (such as B-cells).</p>	<p>3T3 fibroblast cells developed through clonal isolation. Cells were differentiated to an adipose-like state before being used in the screen. See Green H and Meuth M., Cell 3: 127-133 (1974), which is herein incorporated by reference in its entirety.</p> <p>Assays for the activation of transcription through the EGR response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate EGR transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the EGR response element that may be used or routinely modified to test EGR response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays</p>	<p>Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Cancer, Autoimmunity, Allergy and Asthma.</p>
-----	---------	------	---	--	--

				disclosed in: Richards JD, et al., J Immunol, 166(6):3855-3864 (2001); Dinkel, A, et al., J Exp Med, 188(12):2215-2224 (1998); and, Newton, JS, et al., Eur J Immunol 1996 Apr;26(4):811-816 (1996), the contents of each of which are herein incorporated by reference in its entirety. Immune cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary epithelial cells that may be used according to these assays include the Raji cell line.	
310	HLWAE11	1258	Activation of transcription through serum response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved	A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune

				<p>in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC).</p> <p>Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example,</p>
--	--	--	--	---	--

					<p>leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and</p>
--	--	--	--	--	---

					asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
310	HLWAE11	1258	IL-10 in Human T-cell 2B9		
310	HLWAE11	1258	Production of VCAM in endothelial cells (such as human umbilical vein endothelial cells (HUVEC))	Assays for measuring expression of VCAM are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate VCAM expression. For example, FMAT may be used to measure the upregulation of cell surface VCAM-1 expression in endothelial cells. Endothelial cells are cells that line blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation. Exemplary endothelial cells that may be used according to these assays include human umbilical vein	Highly preferred indications include inflammation (acute and chronic), restenosis, atherosclerosis, asthma and allergy. Highly preferred indications include inflammation and inflammatory disorders, immunological disorders, neoplastic disorders (e.g. cancer/tumorigenesis), and cardiovascular disorders (such as described below under "Immune Activity", "Blood-Related Disorders", "Hyperproliferative Disorders" and/or "Cardiovascular Disorders"). Highly preferred indications include neoplasms and cancers such as, for example, leukemia, lymphoma, melanoma, renal cell carcinoma, and prostate, breast, lung, colon, pancreatic,

				endothelial cells (HUVEC), which are available from commercial sources. The expression of VCAM (CD106), a membrane-associated protein, can be upregulated by cytokines or other factors, and contributes to the extravasation of lymphocytes, leucocytes and other immune cells from blood vessels; thus VCAM expression plays a role in promoting immune and inflammatory responses.	esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
310	HLWAE11	1258	Activation of transcription through NFKB response element in immune cells (such as natural killer cells).	Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFKB response element that	Highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), and immunodeficiencies (e.g., as

				<p>may be used or routinely modified to test NFKB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Valle Blazquez et al, Immunology 90(3):455-460 (1997); Aramburau et al., J Exp Med 82(3):801-810 (1995); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. NK cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human NK cells that may be used according to these assays include the NKL cell line, which is a human natural killer cell line established from the peripheral</p>	<p>described below). An additional highly preferred indication is infection (e.g., AIDS, and/or an infectious disease as described below under "Infectious Disease"). Highly preferred indications include neoplastic diseases (e.g., melanoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, melanoma, renal cell carcinoma, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute</p>
--	--	--	--	---	---

				blood of a patient with large granular lymphocytic leukemia. This IL-2 dependent suspension culture cell line has a morphology resembling that of activated NK cells.	lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, suppression of immune reactions to transplanted organs, asthma and allergy.
310	HLWAE11	1258	IL-8 in SW480		
310	HLWAE11	1258	Calcium flux in immune cells (such as monocytes)	Assays for measuring calcium flux are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mobilize calcium. Cells normally have very low concentrations of cytosolic calcium compared to much higher extracellular calcium. Extracellular factors can cause an influx of calcium, leading to activation of calcium responsive signaling	Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of Infection, Inflammation, Atherosclerosis, Hypersensitivity, and Leukemias

				<p>pathways and alterations in cell functions. Exemplary assays that may be used or routinely modified to measure calcium flux in immune cells (such as monocytes) include assays disclosed in: Chan, CC, et al., <i>J Pharmacol Exp Ther</i>, 269(3):891-896 (1994); Andersson, K, et al., <i>Cytokine</i>, 12(12):1784-1787 (2000); Scully, SP, et al., <i>J Clin Invest</i>, 74(2) 589-599 (1984); and, Sullivan, E, et al., <i>Methods Mol Biol</i>, 114:125-133 (1999), the contents of each of which is herein incorporated by reference in its entirety. Cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary cells that may be used according to these assays include the THP-1 monocyte cell line.</p>	
	HLWAO22	1259	Production of MCP-1	<p>MCP-1 FMAT. Assays for immunomodulatory proteins that are produced by a large variety of cells and act to</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing)</p>
311					

				<p>induce chemotaxis and activation of monocytes and T cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, induce chemotaxis, and modulate immune cell activation. Exemplary assays that test for immunomodulatory proteins evaluate the production of cell surface markers, such as monocyte chemoattractant protein (MCP), and the activation of monocytes and T cells. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical</p>	<p>MCP-1 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) MCP-1 production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia,</p>
--	--	--	--	--	---

				<p>approach" Chapter 6:138-160 (2000); Satthaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety.</p> <p>Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art.</p> <p>Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis (bacterial and viral), Lyme Disease, asthma, and allergy Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic</p>
--	--	--	--	---	--

					conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
311	HLWAO22	1259	IL-10 in Human T-cell 2B9		
311	HLWAO22	1259	Activation of transcription through NFAT response in immune cells (such as T-cells).	Assays for the activation of transcription through the Nuclear Factor of Activated T cells (NFAT) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFAT transcription factors and modulate expression of genes involved in immunomodulatory functions. Exemplary assays for transcription through the NFAT response element that may be used or routinely modified to test NFAT-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene	Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred

				<p>66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Serfling et al., Biochim Biophys Acta 1498(1):1-18 (2000); De Boer et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Fraser et al., Eur J Immunol 29(3):838-844 (1999); and Yeseen et al., J Biol Chem 268(19):14285-14293 (1993), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these assays include the JURKAT cell line, which is a suspension culture of leukemia cells that produce IL-2 when stimulated.</p>	<p>indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted</p>
--	--	--	--	---	---

311	HLWAO22	1259	Activation of transcription through serum response element in immune cells (such as natural killer cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate serum response factors and modulate the expression of genes involved in growth and upregulate the function of growth-related genes in many cell types. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and	organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy. A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and
-----	---------	------	--	--	---

				<p>Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Benson et al., J Immunol 153(9):3862-3873 (1994); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary T cells that may be used according to these assays include the NK-YT cell line, which is a human natural killer cell line with cytolytic and cytotoxic activity.</p>	<p>suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
--	--	--	--	--	---

312	HLWAY54	1260	Production of MCP-1	MCP-1 FMAT. Assays for immunomodulatory proteins that are produced by a large variety of cells and act to induce chemotaxis and activation of monocytes and T cells are well known in the art and may be used or routinely	<p>Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p> <p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) MCP-1 production. An alternative highly preferred embodiment of the invention includes a method for</p>
-----	---------	------	---------------------	--	---

				<p>modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, induce chemotaxis, and modulate immune cell activation. Exemplary assays that test for immunomodulatory proteins evaluate the production of cell surface markers, such as monocyte chemoattractant protein (MCP), and the activation of monocytes and T cells. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Satthaporn and Eremin, J R Coll Surg Ednb 45(1):9-19 (2001); and</p>	<p>inhibiting (e.g., reducing) MCP-1 production. A highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Additional highly preferred indications include inflammation and inflammatory disorders. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma,</p>
--	--	--	--	--	---

				<p>Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety.</p> <p>Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art.</p> <p>Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis (bacterial and viral), Lyme Disease, asthma, and allergy Preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p> <p>Highly preferred indications</p>
HLWAY54	1260	Activation of JNK	Kinase assay. JNK kinase		

312	Signaling Pathway in immune cells (such as eosinophils).	assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK kinase activity that may be used or routinely modified to test JNK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which	include asthma, allergy, hypersensitivity reactions, inflammation, and inflammatory disorders. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting or inhibiting immune cell proliferation. Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include boosting an eosinophil-mediated immune response, and suppressing an eosinophil-mediated immune response.
-----	--	---	--

				<p>are herein incorporated by reference in its entirety. Exemplary cells that may be used according to these assays include eosinophils. Eosinophils are important in the late stage of allergic reactions; they are recruited to tissues and mediate the inflammatory response of late stage allergic reaction. Moreover, exemplary assays that may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate signal transduction, cell proliferation, activation, or apoptosis in eosinophils include assays disclosed and/or cited in: Zhang JP, et al., "Role of caspases in dexamethasone-induced apoptosis and activation of c-Jun NH2-terminal kinase and p38 mitogen-activated protein kinase in human eosinophils" Clin Exp Immunol; Oct;122(1):20-7 (2000);</p>	
--	--	--	--	--	--

					Hebestreit H, et al., "Disruption of fas receptor signaling by nitric oxide in eosinophils" J Exp Med; Feb 2;187(3):415-25 (1998); J Allergy Clin Immunol 1999 Sep;104(3 Pt 1):565-74; and, Sousa AR, et al., "In vivo resistance to corticosteroids in bronchial asthma is associated with enhanced phosphorylation of JUN N-terminal kinase and failure of prednisolone to inhibit JUN N-terminal kinase phosphorylation" J Allergy Clin Immunol; Sep;104(3 Pt 1):565-74 (1999); the contents of each of which are herein incorporated by reference in its entirety.	
312	HLWAY54	1260	SEAP in HepG2/Squalene synthetase(stimulation)			
312	HLWAY54	1260	CD152 in Human T cells			
312	HLWAY54	1260	SEAP in OE-33			
312	HLWAY54	1260	SEAP in Senescence Assay			

313	HLWBH18	1261	Inhibition of squalene synthetase gene transcription.	Reporter Assay: construct contains regulatory and coding sequence of squalene synthetase, the first specific enzyme in the cholesterol biosynthetic pathway. See Jiang, et al., J. Biol. Chem. 268:12818-12824(1993), the contents of which are herein incorporated by reference in its entirety. Cells were treated with SID supernatants, and SEAP activity was measured after 72 hours. HepG2 is a human hepatocellular carcinoma cell line (ATCC HB-8065). See Knowles et al., Science. 209:497-9 (1980), the contents of which are herein incorporated by reference in its entirety.	A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell growth. A highly preferred embodiment of the invention
313	HLWBH18	1261	Activation of Endothelial Cell p38 or JNK Signaling Pathway.	Kinase assay. JNK and p38 kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or	

				<p>antagonists of the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK and p38 kinase activity that may be used or routinely modified to test JNK and p38 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Endothelial cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary endothelial cells that may be used according to</p>	<p>includes a method for stimulating endothelial cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting endothelial cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) endothelial cell activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) the activation of and/or inactivating endothelial cells. A highly preferred embodiment of the invention includes a method for</p>
--	--	--	--	---	---

				<p>these assays include human umbilical vein endothelial cells (HUVEC), which are endothelial cells which line venous blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	<p>stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension, aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or</p>
--	--	--	--	---	---

					<p>as described below under "Cardiovascular Disorders").</p> <p>Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization.</p> <p>Highly preferred indications include antiangiogenic activity to treat solid tumors, leukemias, and Kaposi's sarcoma, and retinal disorders.</p> <p>Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma,</p>
--	--	--	--	--	---

					<p>angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly</p>
--	--	--	--	--	--

					<p>preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular degeneration, and treatment /prevention of endometriosis and related conditions. Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as</p>
--	--	--	--	--	--

					described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.
313	HLWBH18	1261	SEAP in OE-33		
314	HLWBI63	1262	CD71 in Human T cells		
315	HLWBK05	1263	Activation of Adipocyte ERK Signaling Pathway	Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely	A highly preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a

				<p>modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that</p>	<p>method for inhibiting adipocyte proliferation. A highly preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) adipocyte activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of (e.g., decreasing) and/or inactivating adipocytes. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"). Highly preferred indications also include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative</p>
--	--	--	--	--	--

				<p>may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>Disorders"). Preferred indications include blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease").</p> <p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below),</p>
--	--	--	--	---	---

					<p>diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyposmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below (particularly of the urinary tract and skin). An</p>
--	--	--	--	--	--

					<p>additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p> <p>Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein.</p> <p>Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Preferred indications include neoplasms and cancer, such as, lymphoma, leukemia and breast, colon, and kidney cancer. Additional preferred indications include melanoma,</p>
--	--	--	--	--	---

316	HLW/BY76	1264	Activation of transcription through serum response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or	prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Highly preferred indications include lipomas and liposarcomas. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
				<p>Assays for the activation of transcription through the Serum Response Element (SRE) are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate the serum response factors and modulate the expression of genes involved in growth. Exemplary assays for transcription through the SRE that may be used or routinely modified to test SRE activity of the polypeptides of the invention (including antibodies and agonists or</p> <p>A preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) TNF alpha production. An alternative preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus,</p>	

				<p>antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); and Black et al., Virus Genes 12(2):105-117 (1997), the content of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that may be used according to these assays include the CTLL cell line, which is an IL-2 dependent suspension culture of T cells with cytotoxic activity.</p>	<p>Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other</p>
--	--	--	--	---	--

316	HLWBY76	1264	CD152 in Human T cells		preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
-----	---------	------	------------------------	--	---

316	HLWBY76	1264	HLA-DR in Human T cells	Kinase assay. Kinase assays, for example an GSK-3 assays, for PI3 kinase signal transduction that regulate glucose metabolism and cell survival are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit glucose metabolism and cell survival. Exemplary assays for PI3 kinase activity that may be used or routinely modified to test PI3 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Nikoulina et al., Diabetes 49(2):263-271 (2000); and Schreyer et al., Diabetes 48(8):1662-1666 (1999), the contents of each of	A highly preferred embodiment of the invention includes a method for increasing adipocyte survival. An alternative highly preferred embodiment of the invention includes a method for decreasing adipocyte survival. A preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"). Preferred indications include
317	HLWCF05	1265	Activation of Adipocyte PI3 Kinase Signalling Pathway		

				<p>which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"), blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other</p>
--	--	--	--	--	---

					<p>diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the</p>
--	--	--	--	--	--

					<p>"Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Highly preferred indications include neoplasms</p>
--	--	--	--	--	---

317	HLWCF05	1265	Activation of JNK Signaling Pathway in immune cells (such as eosinophils).	Kinase assay. JNK kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK kinase activity that may be used or routinely modified to test JNK kinase-induced	and cancer, such as, lipoma, liposarcoma, lymphoma, leukemia and breast, colon, and kidney cancer. Additional highly preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications include asthma, allergy, hypersensitivity reactions, inflammation, and inflammatory disorders. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as
-----	---------	------	--	---	---

				<p>activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety. Exemplary cells that may be used according to these assays include eosinophils. Eosinophils are important in the late stage of allergic reactions; they are recruited to tissues and mediate the inflammatory response of late stage allergic reaction. Moreover, exemplary assays that may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies</p>	<p>described below). Highly preferred indications also include boosting or inhibiting immune cell proliferation. Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include boosting an eosinophil-mediated immune response, and suppressing an eosinophil-mediated immune response.</p>
--	--	--	--	--	--

				<p>and agonists or antagonists of the invention) to modulate signal transduction, cell proliferation, activation, or apoptosis in eosinophils include assays disclosed and/or cited in: Zhang JP, et al., "Role of caspases in dexamethasone-induced apoptosis and activation of c-Jun NH2-terminal kinase and p38 mitogen-activated protein kinase in human eosinophils" Clin Exp Immunol; Oct;122(1):20-7 (2000); Hebestreit H, et al., "Disruption of fas receptor signaling by nitric oxide in eosinophils" J Exp Med; Feb 2;187(3):415-25 (1998); J Allergy Clin Immunol 1999 Sep;104(3 Pt 1):565-74; and, Sousa AR, et al., "In vivo resistance to corticosteroids in bronchial asthma is associated with enhanced phosphorylation of JUN N-terminal kinase and failure of prednisolone to inhibit JUN N-terminal kinase phosphorylation" J Allergy</p>
--	--	--	--	---

					Clin Immunol; Sep;104(3 Pt 1):565-74 (1999); the contents of each of which are herein incorporated by reference in its entirety.	
317	HLWCF05	1265		SEAP in OE-21		
317	HLWCF05	1265		SEAP in OE-33		
317	HLWCF05	1265		Activation of transcription through API response element in immune cells (such as T-cells).	Assays for the activation of transcription through the API response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate growth and other cell functions. Exemplary assays for transcription through the API response element that may be used or routinely modified to test API-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and	Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), and infection (e.g., an infectious disease as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications

				<p>Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety.</p> <p>Human T cells that may be used according to these assays are publicly available (e.g., through the ATCC).</p> <p>Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is an IL-2 and IL-4 responsive suspension-culture cell line.</p>	<p>include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease,</p>
--	--	--	--	--	---

317	HL WCF05	1265	Activation of transcription through CD28 response element in immune cells (such as T-cells).	Assays for the activation of transcription through the CD28 response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate IL-2 expression in T cells. Exemplary assays for transcription through the CD28 response element that may be used or routinely modified to test CD28-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988);	sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease. A highly preferred embodiment of the invention includes a method for stimulating T cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting T cell proliferation. A highly preferred embodiment of the invention includes a method for activating T cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating T cells. A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-2 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-2 production.
-----	----------	------	--	---	--

				<p>McGuire and Iacobelli, J Immunol 159(3):1319-1327 (1997); Parra et al., J Immunol 166(4):2437-2443 (2001); and Butscher et al., J Biol Chem 3(1):552-560 (1998), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC).</p> <p>Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p>	<p>Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Highly preferred indications include neoplastic diseases (e.g., melanoma, renal cell carcinoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, for example, melanoma (e.g., metastatic melanoma), renal cell carcinoma (e.g., metastatic renal cell carcinoma), leukemia, lymphoma (e.g., T cell lymphoma), and prostate,</p>
--	--	--	--	--	---

					breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. A highly preferred indication includes infection (e.g., AIDS, tuberculosis, infections associated with granulomatous disease, and osteoporosis, and/or as described below under "Infectious Disease"). A highly preferred indication is AIDS. Additional highly preferred indications include suppression of immune reactions to transplanted organs and/or tissues, uveitis, psoriasis, and tropical spastic paraparesis. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Preferred indications also
--	--	--	--	--	--

317	HLWCF05	1265	Activation of transcription through NFAT response element in immune cells (such as T-cells).	Assays for the activation of transcription through the Nuclear Factor of Activated T cells (NFAT) response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFAT transcription factors and modulate expression of genes involved in immunomodulatory functions. Exemplary assays for transcription through the	include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.
				Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune	

				<p>NFAT response element that may be used or routinely modified to test NFAT-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Serfling et al., Biochim Biophys Acta 1498(1):1-18 (2000); De Boer et al., Int J Biochem Cell Biol 31(10):1221-1236 (1999); Fraser et al., Eur J Immunol 29(3):838-844 (1999); and Yeseen et al., J Biol Chem 268(19):14285-14293 (1993), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary human T cells that may be used according to these</p>	<p>response. Additional highly preferred indications include inflammation and inflammatory disorders. An additional highly preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease"). Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute</p>
--	--	--	--	--	---

				assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.	lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.
317	HLWCF05	1265	Activation of transcription through NFKB response element in immune cells (such as T-cells).	Assays for the activation of transcription through the NFKB response element are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate NFKB transcription factors and modulate expression of immunomodulatory genes. Exemplary assays for transcription through the NFKB response element that may be used or routinely	Highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), and immunodeficiencies (e.g., as described below). An

			<p>modified to test NFkB-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1998); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Black et al., Virus Gnes 15(2):105-117 (1997); and Fraser et al., 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. T cells that may be used according to these assays are publicly available (e.g., through the ATCC).</p> <p>Exemplary human T cells that may be used according to these assays include the SUPT cell line, which is a suspension culture of IL-2 and IL-4 responsive T cells.</p>	<p>additional highly preferred indication is infection (e.g., AIDS, and/or an infectious disease as described below under "Infectious Disease"). Highly preferred indications include neoplastic diseases (e.g., melanoma, leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, melanoma, renal cell carcinoma, leukemia, lymphoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications also include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL),</p>
--	--	--	---	---

						plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, suppression of immune reactions to transplanted organs, asthma and allergy.
318	HLYAC95	1266	Glucose Production in H4IIE			
318	HLYAC95	1266	Production of IFNgamma using a T cells	IFNgamma FMAT. IFNγ plays a central role in the immune system and is considered to be a proinflammatory cytokine. IFNγ promotes TH1 and inhibits TH2 differentiation; promotes IgG2a and inhibits IgE secretion; induces macrophage activation; and increases MHC expression. Assays for immunomodulatory proteins produced by T cells and NK cells that regulate a variety of inflammatory activities and inhibit TH2 helper cell functions are well	A highly preferred embodiment of the invention includes a method for stimulating the production of IFNγ. An alternative highly preferred embodiment of the invention includes a method for inhibiting the production of IFNγ. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., viral	

				<p>known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, regulate inflammatory activities, modulate TH2 helper cell function, and/or mediate humoral or cell-mediated immunity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as Interferon gamma (IFNg), and the activation of T cells. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204 (1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160</p>	<p>infections, tuberculosis, infections associated with chronic granulomatous disease and malignant osteoporosis, and/or as described below under "Infectious Disease"). Highly preferred indications include autoimmune disease (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below), immunodeficiency (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders. Additional preferred indications include idiopathic pulmonary fibrosis. Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred</p>
--	--	--	--	---	---

			<p>(2000); Gonzalez et al., J Clin Lab Anal 8(5):225-233 (1995); Billiau et al., Ann NY Acad Sci 856:22-32 (1998); Boehm et al., Annu Rev Immunol 15:749-795 (1997), and Rheumatology (Oxford) 38(3):214-20 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human T cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human T cells are primary human lymphocytes that mature in the thymus and express a T Cell receptor and CD3, CD4, or CD8. These cells mediate humoral or cell-mediated immunity and may be preactivated to enhance responsiveness to immunomodulatory factors.</p>	<p>indications include neoplasms and cancers, such as, for example, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, asthma and allergy.</p>
--	--	--	--	---

318	HL YAC95	1266	Stimulation of insulin secretion from pancreatic beta cells.	<p>Assays for measuring secretion of insulin are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate insulin secretion. For example, insulin secretion is measured by FMAT using anti-rat insulin antibodies. Insulin secretion from pancreatic beta cells is upregulated by glucose and also by certain proteins/peptides, and dysregulation is a key component in diabetes. Exemplary assays that may be used or routinely modified to test for stimulation of insulin secretion (from pancreatic cells) by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Ahren, B., et al., Am J Physiol, 277(4 Pt 2):R959-66 (1999); Li, M., et al., Endocrinology,</p>	<p>A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hypoglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia,</p>
-----	----------	------	--	--	--

				<p>138(9):3735-40 (1997); Kim, K.H., et al., FEBS Lett, 377(2):237-9 (1995); and, Miraglia S et. al., Journal of Biomolecular Screening, 4:193-204 (1999), the contents of each of which is herein incorporated by reference in its entirety. Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include rat INS-1 cells. INS-1 cells are a semi-adherent cell line established from cells isolated from an X-ray induced rat transplantable insulinoma. These cells retain characteristics typical of native pancreatic beta cells including glucose inducible insulin secretion. References: Asfari et al. Endocrinology 1992 130:167.</p>	<p>endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
318	HL YAC95	1266	Hexosaminidase in RBL-2H3		
	HL YAF80	1267	SEAP in		

319			NK16/STAT6			
320	HL YAN59	1268	CD152 in Human T cells			
320	HL YAN59	1268	HLA-DR in Human T cells			
320	HL YAN59	1268	Production of VCAM in endothelial cells (such as human umbilical vein endothelial cells (HUV EC))	Assays for measuring expression of VCAM are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate VCAM expression. For example, FMAT may be used to measure the upregulation of cell surface VCAM-1 expression in endothelial cells. Endothelial cells are cells that line blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation. Exemplary endothelial cells that may be used according to these assays include human umbilical vein endothelial cells (HUV EC), which are available from	Highly preferred indications include inflammation (acute and chronic), restnosis, atherosclerosis, asthma and allergy. Highly preferred indications include inflammation and inflammatory disorders, immunological disorders, neoplastic disorders (e.g. cancer/tumorigenesis), and cardiovascular disorders (such as described below under "Immune Activity", "Blood-Related Disorders", "Hyperproliferative Disorders" and/or "Cardiovascular Disorders"). Highly preferred indications include neoplasms and cancers such as, for example, leukemia, lymphoma, melanoma, renal cell carcinoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other	

				commercial sources. The expression of VCAM (CD106), a membrane-associated protein, can be upregulated by cytokines or other factors, and contributes to the extravasation of lymphocytes, leucocytes and other immune cells from blood vessels; thus VCAM expression plays a role in promoting immune and inflammatory responses.	preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
321	HL YAP91	1269	Activation of Adipocyte ERK Signaling Pathway	Kinase assay. Kinase assays, for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced	A highly preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte proliferation. A highly preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. A highly

				<p>activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Le Marchand-Brustel Y, Exp Clin Endocrinol Diabetes 107(2):126-132 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein incorporated by reference in its entirety.</p> <p>Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed</p>	<p>preferred embodiment of the invention includes a method for stimulating (e.g., increasing) adipocyte activation. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of (e.g., decreasing) and/or inactivating adipocytes. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders").</p> <p>Highly preferred indications also include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"). Preferred indications include blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders</p>
--	--	--	--	---	--

				through clonal isolation and undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.	(e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma,
--	--	--	--	--	--

					cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below (particularly of the urinary tract and skin). An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance. Additional highly preferred
--	--	--	--	--	---

					<p>indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein.</p> <p>Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Preferred indications include neoplasms and cancer, such as, lymphoma, leukemia and breast, colon, and kidney cancer. Additional preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer.</p> <p>Highly preferred indications include lipomas and liposarcomas. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.</p>
--	--	--	--	--	--

322	HL YAZ61	1270	<p>Activation of JNK Signaling Pathway in immune cells (such as eosinophils).</p>	<p>Kinase assay. JNK kinase assays for signal transduction that regulate cell proliferation, activation, or apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and apoptosis. Exemplary assays for JNK kinase activity that may be used or routinely modified to test JNK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Gupta et al., Exp Cell Res 247(2): 495-504 (1999); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999);</p>	<p>Highly preferred indications include asthma, allergy, hypersensitivity reactions, inflammation, and inflammatory disorders. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting or inhibiting immune cell proliferation. Preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include boosting an eosinophil-mediated immune response, and suppressing an eosinophil-mediated immune</p>
-----	----------	------	---	---	--

				<p>the contents of each of which are herein incorporated by reference in its entirety. Exemplary cells that may be used according to these assays include eosinophils. Eosinophils are important in the late stage of allergic reactions; they are recruited to tissues and mediate the inflammatory response of late stage allergic reaction. Moreover, exemplary assays that may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to modulate signal transduction, cell proliferation, activation, or apoptosis in eosinophils include assays disclosed and/or cited in: Zhang JP, et al., "Role of caspases in dexamethasone-induced apoptosis and activation of c-Jun NH2-terminal kinase and p38 mitogen-activated protein kinase in human eosinophils" Clin Exp Immunol;</p>	response.
--	--	--	--	--	-----------

					Oct;122(1):20-7 (2000); Hebestreit H, et al., "Disruption of fas receptor signaling by nitric oxide in eosinophils" J Exp Med; Feb 2;187(3):415-25 (1998); J Allergy Clin Immunol 1999 Sep;104(3 Pt 1):565-74; and, Sousa AR, et al., "In vivo resistance to corticosteroids in bronchial asthma is associated with enhanced phosphorylation of JUN N-terminal kinase and failure of prednisolone to inhibit JUN N-terminal kinase phosphorylation" J Allergy Clin Immunol; Sep;104(3 Pt 1):565-74 (1999); the contents of each of which are herein incorporated by reference in its entirety.	
323	HL YBD32	1271	Activation of transcription through AP1 response element in immune cells (such as T-cells).	Assays for the activation of transcription through the AP1 response element are known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to	Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related	

				<p>modulate growth and other cell functions. Exemplary assays for transcription through the AP1 response element that may be used or routinely modified to test AP1-response element activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Berger et al., Gene 66:1-10 (1988); Cullen and Malm, Methods in Enzymol 216:362-368 (1992); Henthorn et al., Proc Natl Acad Sci USA 85:6342-6346 (1988); Rellahan et al., J Biol Chem 272(49):30806-30811 (1997); Chang et al., Mol Cell Biol 18(9):4986-4993 (1998); and Fraser et al., Eur J Immunol 29(3):838-844 (1999), the contents of each of which are herein incorporated by reference in its entirety. Mouse T cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse T cells that</p>	<p>Disorders”), and infection (e.g., an infectious disease as described below under “Infectious Disease”). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional highly preferred indications include inflammation and inflammatory disorders. Highly preferred indications also include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under “Hyperproliferative Disorders”). Highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic</p>
--	--	--	--	--	--

				may be used according to these assays include the HT2 cell line, which is an IL-2 dependent suspension culture cell line that also responds to IL-4.	conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include arthritis, asthma, AIDS, allergy, anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, granulomatous disease, inflammatory bowel disease, sepsis, psoriasis, suppression of immune reactions to transplanted organs and tissues, endocarditis, meningitis, and Lyme Disease.
324	HL YES38	1272	Activation of Adipocyte PI3 Kinase Signalling Pathway	Kinase assay. Kinase assays, for example an GSK-3 assays, for PI3 kinase signal transduction that regulate glucose metabolism and cell survival are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit glucose	<p>A highly preferred embodiment of the invention includes a method for increasing adipocyte survival</p> <p>An alternative highly preferred embodiment of the invention includes a method for decreasing adipocyte survival.</p> <p>A preferred embodiment of the invention includes a method for stimulating adipocyte proliferation. An alternative highly preferred embodiment</p>

				<p>metabolism and cell survival. Exemplary assays for PI3 kinase activity that may be used or routinely modified to test PI3 kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Nikoulina et al., Diabetes 49(2):263-271 (2000); and Schreyer et al., Diabetes 48(8):1662-1666 (1999), the contents of each of which are herein incorporated by reference in its entirety. Mouse adipocyte cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary mouse adipocyte cells that may be used according to these assays include 3T3-L1 cells. 3T3-L1 is an adherent mouse preadipocyte cell line that is a continuous substrain of 3T3 fibroblast cells developed through clonal isolation and</p>	<p>of the invention includes a method for inhibiting adipocyte proliferation. A preferred embodiment of the invention includes a method for stimulating adipocyte differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting adipocyte differentiation. Highly preferred indications include endocrine disorders (e.g., as described below under "Endocrine Disorders"). Preferred indications include neoplastic diseases (e.g., lipomas, liposarcomas, and/or as described below under "Hyperproliferative Disorders"), blood disorders (e.g., hypertension, congestive heart failure, blood vessel blockage, heart disease, stroke, impotence and/or as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune</p>
--	--	--	--	--	--

				<p>undergo a pre-adipocyte to adipose-like conversion under appropriate differentiation conditions known in the art.</p>	<p>Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g.,</p>
--	--	--	--	--	--

					<p>heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the</p> <p>"Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture).</p> <p>An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
--	--	--	--	--	---

					Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein. Additional highly preferred indications include, hypertension, coronary artery disease, dyslipidemia, gallstones, osteoarthritis, degenerative arthritis, eating disorders, fibrosis, cachexia, and kidney diseases or disorders. Highly preferred indications include neoplasms and cancer, such as, lipoma, liposarcoma, lymphoma, leukemia and breast, colon, and kidney cancer. Additional highly preferred indications include melanoma, prostate, lung, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
	HLYES38	1272	IFNg in Human T-		

			<p>stimulating apoptosis of endothelial cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing) apoptosis of endothelial cells. A highly preferred embodiment of the invention includes a method for stimulating angiogenesis. An alternative highly preferred embodiment of the invention includes a method for inhibiting angiogenesis. A highly preferred embodiment of the invention includes a method for reducing cardiac hypertrophy. An alternative highly preferred embodiment of the invention includes a method for inducing cardiac hypertrophy. Highly preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), and disorders of the cardiovascular system (e.g., heart disease, congestive heart failure, hypertension,</p>
		<p>Endothelial cells that may be used according to these assays are publicly available (e.g., through commercial sources). Exemplary endothelial cells that may be used according to these assays include bovine aortic endothelial cells (bAEC), which are an example of endothelial cells which line blood vessels and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation.</p>	

					<p>aortic stenosis, cardiomyopathy, valvular regurgitation, left ventricular dysfunction, atherosclerosis and atherosclerotic vascular disease, diabetic nephropathy, intracardiac shunt, cardiac hypertrophy, myocardial infarction, chronic hemodynamic overload, and/or as described below under “Cardiovascular Disorders”).</p> <p>Highly preferred indications include cardiovascular, endothelial and/or angiogenic disorders (e.g., systemic disorders that affect vessels such as diabetes mellitus, as well as diseases of the vessels themselves, such as of the arteries, capillaries, veins and/or lymphatics). Highly preferred are indications that stimulate angiogenesis and/or cardiovascularization. Highly preferred are indications that inhibit angiogenesis and/or cardiovascularization.</p> <p>Highly preferred indications include antiangiogenic activity to treat solid tumors,</p>
--	--	--	--	--	---

					<p>leukemias, and Kaposi's sarcoma, and retinal disorders. Highly preferred indications include neoplasms and cancer, such as, Kaposi's sarcoma, hemangioma (capillary and cavernous), glomus tumors, telangiectasia, bacillary angiomatosis, hemangioendothelioma, angiosarcoma, haemangiopericytoma, lymphangioma, lymphangiosarcoma. Highly preferred indications also include cancers such as, prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver, and urinary cancer. Preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Highly preferred indications also include arterial disease, such as, atherosclerosis, hypertension, coronary artery disease, inflammatory vasculitides, Reynaud's</p>
--	--	--	--	--	--

					<p>disease and Reynaud's phenomenon, aneurysms, restenosis; venous and lymphatic disorders such as thrombophlebitis, lymphangitis, and lymphedema; and other vascular disorders such as peripheral vascular disease, and cancer. Highly preferred indications also include trauma such as wounds, burns, and injured tissue (e.g., vascular injury such as, injury resulting from balloon angioplasty, and atherosclerotic lesions), implant fixation, scarring, ischemia reperfusion injury, rheumatoid arthritis, cerebrovascular disease, renal diseases such as acute renal failure, and osteoporosis. Additional highly preferred indications include stroke, graft rejection, diabetic or other retinopathies, thrombotic and coagulative disorders, vasculitis, lymph angiogenesis, sexual disorders, age-related macular</p>
--	--	--	--	--	--

	HMADS41	1273	Activation of	Kinase assay. Kinase assays,	<p>degeneration, and treatment /prevention of endometriosis and related conditions.</p> <p>Additional highly preferred indications include fibromas, heart disease, cardiac arrest, heart valve disease, and vascular disease. Preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders").</p> <p>Preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and immunodeficiencies (e.g., as described below). Additional preferred indications include inflammation and inflammatory disorders (such as acute and chronic inflammatory diseases, e.g., inflammatory bowel disease and Crohn's disease), and pain management.</p>
--	---------	------	---------------	------------------------------	--

325	Hepatocyte ERK Signaling Pathway	<p>for example an Elk-1 kinase assay, for ERK signal transduction that regulate cell proliferation or differentiation are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to promote or inhibit cell proliferation, activation, and differentiation. Exemplary assays for ERK kinase activity that may be used or routinely modified to test ERK kinase-induced activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in Forrer et al., Biol Chem 379(8-9):1101-1110 (1998); Kyriakis JM, Biochem Soc Symp 64:29-48 (1999); Chang and Karin, Nature 410(6824):37-40 (2001); and Cobb MH, Prog Biophys Mol Biol 71(3-4):479-500 (1999); the contents of each of which are herein</p>	<p>embodiment of the invention includes a method for stimulating hepatocyte cell proliferation. An alternative highly preferred embodiment of the invention includes a method for inhibiting hepatocyte cell proliferation. A highly preferred embodiment of the invention includes a method for stimulating hepatocyte cell differentiation. An alternative highly preferred embodiment of the invention includes a method for inhibiting hepatocyte cell differentiation. A highly preferred embodiment of the invention includes a method for activating hepatocyte cells. An alternative highly preferred embodiment of the invention includes a method for inhibiting the activation of and/or inactivating hepatocyte cells. Highly preferred indications include disorders of the liver and/or endocrine disorders (e.g., as described below under "Endocrine</p>
-----	----------------------------------	---	--

				<p>incorporated by reference in its entirety. Rat liver hepatoma cells that may be used according to these assays are publicly available (e.g., through the ATCC). Exemplary rat liver hepatoma cells that may be used according to these assays include H4Ile cells, which are known to respond to glucocorticoids, insulin, or cAMP derivatives.</p>	<p>Disorders"). Preferred indications include neoplastic diseases (e.g., as described below under "Hyperproliferative Disorders"), blood disorders (e.g., as described below under "Immune Activity", "Cardiovascular Disorders", and/or "Blood-Related Disorders"), immune disorders (e.g., as described below under "Immune Activity"), neural disorders (e.g., as described below under "Neural Activity and Neurological Diseases"), and infection (e.g., as described below under "Infectious Disease"). A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below),</p>
--	--	--	--	--	--

					<p>diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g., heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal</p>
--	--	--	--	--	--

					<p>tunnel syndrome and Dupuytren's contracture).</p> <p>An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p> <p>Additional highly preferred indications are disorders of the musculoskeletal systems including myopathies, muscular dystrophy, and/or as described herein.</p> <p>Additional highly preferred indications include, hepatitis, jaundice, gallstones, cirrhosis of the liver, degenerative or necrotic liver disease, alcoholic liver diseases, fibrosis, liver regeneration, metabolic disease, dyslipidemia and cholesterol metabolism.</p> <p>Additional highly preferred indications include neoplasms and cancers, such as,</p>
--	--	--	--	--	---

325	HMADS41	1273	Regulation of apoptosis of immune cells (such as mast cells).	Caspase Apoptosis. Assays for caspase apoptosis are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate caspase protease-mediated apoptosis in immune cells (such as, for example, in mast cells). Mast cells are found in connective and mucosal tissues throughout the body, and their activation via immunoglobulin E - antigen, promoted by T helper cell type 2 cytokines, is an	hepatocarcinomas, other liver cancers, and colon and pancreatic cancer. Preferred indications also include prostate, breast, lung, esophageal, stomach, brain, and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia.
					Preferred embodiments of the invention include using polypeptides of the invention (or antibodies, agonists, or antagonists thereof) in detection, diagnosis, prevention, and/or treatment of asthma, allergy, hypersensitivity and inflammation.

				<p>important component of allergic disease. Dysregulation of mast cell apoptosis may play a role in allergic disease and mast cell tumor survival. Exemplary assays for caspase apoptosis that may be used or routinely modified to test caspase apoptosis activity induced by polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include the assays disclosed in: Masuda A, et al., J Biol Chem, 276(28):26107-26113 (2001); Yeatman CF 2nd, et al., J Exp Med, 192(8):1093-1103 (2000); Lee et al., FEBS Lett 485(2-3): 122-126 (2000); Nor et al., J Vasc Res 37(3): 209-218 (2000); and Karsan and Harlan, J Atheroscler Thromb 3(2): 75-80 (1996); the contents of each of which are herein incorporated by reference in its entirety. Immune cells that may be used according to these assays are publicly available (e.g., through commercial sources).</p>	
--	--	--	--	---	--

				Exemplary immune cells that may be used according to these assays include mast cells such as the HMC human mast cell line.	
326	HMADU73	1274	Production of IL-6	<p>IL-6 FMAT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described</p>

				<p>invention) to mediate immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may</p>	<p>below) and immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred</p>
--	--	--	--	---	---

			<p>be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").</p>
326	HMADU73	1274	<p>Production of TNF alpha by dendritic cells</p>	<p>TNFα FMAT. Assays for immunomodulatory proteins produced by activated macrophages, T cells,</p> <p>A highly preferred embodiment of the invention includes a method for inhibiting (e.g., decreasing)</p>

				<p>fibroblasts, smooth muscle, and other cell types that exert a wide variety of inflammatory and cytotoxic effects on a variety of cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate immunomodulation, modulate inflammation and cytotoxicity. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines such as tumor necrosis factor alpha (TNFa), and the induction or inhibition of an inflammatory or cytotoxic response. Such assays that may be used or routinely modified to test immunomodulatory activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-</p>	<p>TNF alpha production. An alternative highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) TNF alpha production. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response. Additional highly preferred indications include inflammation and inflammatory disorders, and treating joint damage in patients with rheumatoid arthritis. An additional highly preferred indication is sepsis.</p>
--	--	--	--	--	---

				<p>204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); Verhasselt et al., Eur J Immunol 28(11):3886-3890 (1998); Dahlen et al., J Immunol 160(7):3585-3593 (1998); Verhasselt et al., J Immunol 158:2919-2925 (1997); and Nardelli et al., J Leukoc Biol 65:822-828 (1999), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.</p>	<p>Highly preferred indications include neoplastic diseases (e.g., leukemia, lymphoma, and/or as described below under "Hyperproliferative Disorders"). Additionally, highly preferred indications include neoplasms and cancers, such as, leukemia, lymphoma, melanoma, glioma (e.g., malignant glioma), solid tumors, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), plasmacytomas, multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, neutropenia,</p>
--	--	--	--	---	---

					neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, Lyme Disease, cardiac reperfusion injury, and asthma and allergy. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
326	HMADU73	1274	IgG in Human B cells SAC		
326	HMADU73	1274	CD152 in Human T cells		
326	HMADU73	1274	Production of IL-10 and activation of T-cells.	Assays for production of IL-10 and activation of T-cells are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to stimulate or inhibit production of IL-10 and/or activation of T-cells. Exemplary assays that may be used or routinely modified to assess the ability of	Highly preferred indications include allergy and asthma. Additional highly preferred indications include immune and hematopoietic disorders (e.g., as described below under "Immune Activity", and "Blood-Related Disorders"), autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), immunodeficiencies (e.g., as

				<p>polypeptides and antibodies of the invention (including agonists or antagonists of the invention) to modulate IL-10 production and/or T-cell proliferation include, for example, assays such as disclosed and/or cited in: Robinson, DS, et al., "Th-2 cytokines in allergic disease" Br Med Bull; 56 (4): 956-968 (2000), and Cohn, et al., "T-helper type 2 cell-directed therapy for asthma" Pharmacology & Therapeutics; 88: 187-196 (2000); the contents of each of which are herein incorporated by reference in their entirety. Exemplary cells that may be used according to these assays include Th2 cells. IL10 secreted from Th2 cells may be measured as a marker of Th2 cell activation. Th2 cells are a class of T cells that secrete IL4, IL10, IL13, IL5 and IL6. Factors that induce differentiation and activation of Th2 cells play a major role in the initiation and</p>	described below), boosting a T cell-mediated immune response, and suppressing a T cell-mediated immune response.
--	--	--	--	--	--

				pathogenesis of allergy and asthma. Primary T helper 2 cells are generated via in vitro culture under Th2 polarizing conditions using peripheral blood lymphocytes isolated from cord blood.	
327	HMAMI15	1275	Stimulation of Calcium Flux in pancreatic beta cells.	Assays for measuring calcium flux are well-known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mobilize calcium. For example, the FLPR assay may be used to measure influx of calcium. Cells normally have very low concentrations of cytosolic calcium compared to much higher extracellular calcium. Extracellular factors can cause an influx of calcium, leading to activation of calcium responsive signaling pathways and alterations in cell functions. Exemplary assays that may be used or routinely modified to measure calcium flux by polypeptides of the	A highly preferred indication is diabetes mellitus. An additional highly preferred indication is a complication associated with diabetes (e.g., diabetic retinopathy, diabetic nephropathy, kidney disease (e.g., renal failure, nephropathy and/or other diseases and disorders as described in the "Renal Disorders" section below), diabetic neuropathy, nerve disease and nerve damage (e.g., due to diabetic neuropathy), blood vessel blockage, heart disease, stroke, impotence (e.g., due to diabetic neuropathy or blood vessel blockage), seizures, mental confusion, drowsiness, nonketotic hyperglycemic-hyperosmolar coma, cardiovascular disease (e.g.,

			<p>invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in: Satin LS, et al., Endocrinology, 136(10):4589-601 (1995); Mogami H, et al., Endocrinology, 136(7):2960-6 (1995); Richardson SB, et al., Biochem J, 288 (Pt 3):847-51 (1992); and, Meats, JE, et al., Cell Calcium 1989 Nov-Dec;10(8):535-41 (1989), the contents of each of which is herein incorporated by reference in its entirety.</p> <p>Pancreatic cells that may be used according to these assays are publicly available (e.g., through the ATCC) and/or may be routinely generated. Exemplary pancreatic cells that may be used according to these assays include HIT15 Cells. HIT15 are an adherent epithelial cell line established from Syrian hamster islet cells transformed with SV40. These cells express glucagon, somatostatin, and glucocorticoid receptors. The cells secrete insulin, which is</p>	<p>heart disease, atherosclerosis, microvascular disease, hypertension, stroke, and other diseases and disorders as described in the "Cardiovascular Disorders" section below), dyslipidemia, endocrine disorders (as described in the "Endocrine Disorders" section below), neuropathy, vision impairment (e.g., diabetic retinopathy and blindness), ulcers and impaired wound healing, and infection (e.g., infectious diseases and disorders as described in the "Infectious Diseases" section below, especially of the urinary tract and skin), carpal tunnel syndrome and Dupuytren's contracture).</p> <p>An additional highly preferred indication is obesity and/or complications associated with obesity. Additional highly preferred indications include weight loss or alternatively, weight gain. Additional highly preferred indications are complications associated with insulin resistance.</p>
--	--	--	--	--

					stimulated by glucose and glucagon and suppressed by somatostatin or glucocorticoids. ATTC# CRL-1777 Refs: Lord and Ashcroft. Biochem. J. 219: 547-551; Santerre et al. Proc. Natl. Acad. Sci. USA 78: 4339-4343, 1981.	
327	HMAMI15	1275	CD152 in Human T cells			
327	HMAMI15	1275	Activation of Transcription		Assays for activation of transcription are well-known in the art and may be used and routinely modified to assess ability of polypeptides of the invention to inhibit or activate transcription. An example of such an assay follows: Cells were pretreated with SID supernatants or controls for 15-18 hours. SEAP activity was measured after 48 hours. LS174T is an epithelial colon adenocarcinoma cell line. Its tumorigenicity in nude mice make cell line LS174T a model for studies on the mechanism of synthesis and secretion of specific tumoral markers in colon cancer. See, Patan et al.,	

				Circ Res, 89(8):732-39 (2001), the contents of which are herein incorporated by reference in its entirety.	
328	HMDAE65	1276	Production of IL-6	<p>IL-6 FMAT. IL-6 is produced by T cells and has strong effects on B cells. IL-6 participates in IL-4 induced IgE production and increases IgA production (IgA plays a role in mucosal immunity). IL-6 induces cytotoxic T cells. Deregulated expression of IL-6 has been linked to autoimmune disease, plasmacytomas, myelomas, and chronic hyperproliferative diseases. Assays for immunomodulatory and differentiation factor proteins produced by a large variety of cells where the expression level is strongly regulated by cytokines, growth factors, and hormones are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to mediate</p>	<p>A highly preferred embodiment of the invention includes a method for stimulating (e.g., increasing) IL-6 production. An alternative highly preferred embodiment of the invention includes a method for inhibiting (e.g., reducing) IL-6 production. A highly preferred indication is the stimulation or enhancement of mucosal immunity. Highly preferred indications include blood disorders (e.g., as described below under "Immune Activity", "Blood-Related Disorders", and/or "Cardiovascular Disorders"), and infection (e.g., as described below under "Infectious Disease"). Highly preferred indications include autoimmune diseases (e.g., rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis and/or as described below) and</p>

				<p>immunomodulation and differentiation and modulate T cell proliferation and function. Exemplary assays that test for immunomodulatory proteins evaluate the production of cytokines, such as IL-6, and the stimulation and upregulation of T cell proliferation and functional activities. Such assays that may be used or routinely modified to test immunomodulatory and differentiation activity of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) include assays disclosed in Miraglia et al., J Biomolecular Screening 4:193-204(1999); Rowland et al., "Lymphocytes: a practical approach" Chapter 6:138-160 (2000); and Verhasselt et al., J Immunol 158:2919-2925 (1997), the contents of each of which are herein incorporated by reference in its entirety. Human dendritic cells that may be used according to these</p>	<p>immunodeficiencies (e.g., as described below). Highly preferred indications also include boosting a B cell-mediated immune response and alternatively suppressing a B cell-mediated immune response. Highly preferred indications include inflammation and inflammatory disorders. Additional highly preferred indications include asthma and allergy. Highly preferred indications include neoplastic diseases (e.g., myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and/or as described below under "Hyperproliferative Disorders"). Highly preferred indications include neoplasms and cancers, such as, myeloma, plasmacytoma, leukemia, lymphoma, melanoma, and prostate, breast, lung, colon, pancreatic, esophageal, stomach, brain, liver and urinary cancer. Other preferred indications include benign</p>
--	--	--	--	--	---

				assays may be isolated using techniques disclosed herein or otherwise known in the art. Human dendritic cells are antigen presenting cells in suspension culture, which, when activated by antigen and/or cytokines, initiate and upregulate T cell proliferation and functional activities.	dysproliferative disorders and pre-neoplastic conditions, such as, for example, hyperplasia, metaplasia, and/or dysplasia. Preferred indications include anemia, pancytopenia, leukopenia, thrombocytopenia, Hodgkin's disease, acute lymphocytic anemia (ALL), multiple myeloma, Burkitt's lymphoma, arthritis, AIDS, granulomatous disease, inflammatory bowel disease, sepsis, neutropenia, neutrophilia, psoriasis, suppression of immune reactions to transplanted organs and tissues, hemophilia, hypercoagulation, diabetes mellitus, endocarditis, meningitis, and Lyme Disease. An additional preferred indication is infection (e.g., an infectious disease as described below under "Infectious Disease").
329	HMDAM24	1277	Protection from Endothelial Cell Apoptosis.	Caspase Apoptosis Rescue. Assays for caspase apoptosis rescue are well known in the art and may be used or routinely modified to assess	A highly preferred embodiment of the invention includes a method for stimulating endothelial cell growth. An alternative highly